

REFERENCES

References

- Abbas, M. S. T.; Hanounik, S. P.; Mousa, S. A. and Al-Bagham, S. H. (2000): Soil application of entomopathogenic nematodes as a new approach for controlling *Rhynchophorus ferrugineus* on date palm. International Journal of Nematology Vol. 10, No. (2): 215-218.
- Abbas, M. S. T.; Shidi, R. H.; Jumah, S. and Al-Khatry, S. A. (2008): Utilization of *Goniozus* sp. (Hym.: Bethylidae) as a bio-control agent against the lesser date moth, *Batrachedra amydraula* (Meyrick) (Lep.: Batrachedridae) in date palm orchards in Sultanate of Oman.
- Abbas, M. S.T. and Hanounik, S. P. (1999): Pathogenicity of entomopathogenic nematodes to the red palm weevil, *Rhynchophorus ferrugineus*. International Journal of Nematology Vol. 9, No. (1): 84-86.
- Abdallah, F. F. and Tadros, A. A. (1994): Biological studies on the date stone beetle, *Coccotrypes dactyliperda* (Fabricius), (Coleoptera : Scolytidae). Alex. Sci. Exch., Vol. 15, No. (1): 1-9.
- Abdallah, F. F. and Al-Khatry, S. A. (2000a): The Effectiveness of trunk injection and fumigation for the control of the red palm weevil, *Rhynchophorus ferrugineus* Oliver in date palm. Journal of Plant Protection in the Tropics, 13 (1): 17-21.
- Abdallah, F. F. and Al-Khatry, S. A. (2000b): Efficacy of different attractant traps on red palm weevil *Rhynchophorus ferrugineus* (Curculionidae: Coleoptera). Proceedings of First Workshop on Control of Date Palm Red Weevil 20-22 November 2000, 24-26 Shabaan 1421H. Ministry of High Education, King Faisal University. Kingdom of Saudi Arabia, 79-87.
- Abdallah, F. F. and AL-Khatry, S. A. (2003): Seasonal abundance of *Oryctes agamemnon* Burmeister, (Coleoptera: Dynastidae) in Sultanate of Oman.
- International Scientific Meeting of the Date Palm, 19-22 Ragab 1424H-16-19 September 2003, College of Agriculture and Veterinary Medicine, King Saud University-Qassim, Ministry of Higher Science, Kingdom of Saudi Arabia.
- Abdallah, F. F. and Al-Khatry, S. A. (2005): The Effect of Pheromone, Kairomone and Food Bait on Attracting adults of Red Palm Weevil *Rhynchophorus ferrugineus* in the Sultanate of Oman in Date Palm Plantations. The Third International Conference of Plant Protection Research Institute, 26-29 November 2005. Egyption Journal of Agricultural Research, 83: 169-177.
- Abdallah, F. F.; Al-Zidjali, T. S. and Al-Khatry, S. A. (1998a): Biology of *Ommatissus lybicus* Bergevin under field and laboratory conditions during spring 1995. Sultan Qaboos University, Proc. IPM, CONF., 23-25 Febrauray 1998, 75-79.
- Abdallah, F. F.; Al-Zidjali, T. S. and Al-Khatry, S. A. (1998b): Biology of *Batrachedra amydraula* Meyrick (Lepidoptera: Cosmopterygidae) under laboratory condition. Sultan Qaboos University, Proc. IPM, CONF., 23-25 Febrauray 1998, 81-86.
- Abdel Qader, H. H. and Al-Husseini, M. S. (1997): Palm diseases (Prevention and control). Kingdom of Saudi Arabia, Riyadh, 130 pp.
- Abdel Salam, A. L. (1993): Insect pests in Egypt and Arab Countries and its Control Measures, Part Two. Insect pests which attack Vegetables, Fruits and Ornamental Plants: 781 pp. (In Arabic).
- Abdelkhalek, L.; Afellah, M. and Smaili, C. (1998): Biology and biological control of *Planococcus citri* R. (Hom., Pseudococcidae) on citrus in the Loukos region of Morocco. Mededelingen-Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen, Universiteit Gent, 63(2b): 483-488.

References

- Abdul Ahad, I. and Jassim, H.K. (1983): The life cycle of *Parlatoria blanchardii* (Targ.) (Diaspididae, Homoptera). Arab Journal of Plant Protection, 1: 22-24.
- Abdul Hag, K. and Akmal, M. (1972): Insect pests of date palm and their control. The Punjab Fruit Journal, 33: 208-212.
- Abdul Wahab, W. (1975): Notes on some natural enemies of Lesser Date Moth (Homera) on date palm and dates, Ministry of Agriculture, Baghdad, Iraq. (In Arabic).
- Abivardi, C. (2001): Iranian entomology: an introduction. Volume 2. Applied entomology. Springer-Verlag, Berlin, Heidelberg, New York, 445-1033.
- Abraham, V. A. (1971): Note on an effective method of preventing entry by red-weevil, *Rhynchophorus ferrugineus* Fabricius (Curculionidae: Coleoptera), into the stem of coconut palm through cut petioles. Indian Journal of Agricultural Sciences, 41 (12):1130-1131.
- Abraham, V. A.; Koya, K. M. A and Kurian, C. (1989): Integrated management of red palm weevil (*Rhynchophorus ferrugineus* F.) in coconut gardens. Journal of Plantation Crops, 16 (Supplement):159-162.
- Abraham, V. A.; Kurian, C. and Nayer, N. M. (1973): *Chelisoches morio* F. (Forficulidae:Dermoptera), a predator on eggs and early instar grubs of the red palm weevil, *Rhynchophorus ferrugineus* F. (Curculionidae: Coleoptera). Journal of Plantation Crops, (1): 147-152.
- Abraham, V.A. and Kurian, C. (1975): An integrated approach to the control of *Rhynchophorus ferrugineus* F. the red weevil of coconut palm. Paper Fourth Session FAO Technical Working Party on Coconut Production, Protection and Processing, Kingston (Jamaica).
- Abu Thuraya, N. H. (1979): A list of date palm pests in Saudi Arabia. Plant Protection Res. Div. Res. Dept., Ministry of Agric. and water, Riyadh.
- Afshari, M.; Mossadegh, S.; Soleyman-Nejadian, E. and Kamali, K. (2007): Geographical distribution and host plants of *Stethorus gilvifrons* (Mulsant) (Col.: Coccinellidae) and its biology under laboratory conditions in Khuzestan province. Journal of Agricultural Science and Natural Resources 14, 201-210.
- Agrios, G. N. (1978): Plant Pathology, Second Edition, Department of Plant Pathology, University of Florida, USA, Elsevier Academic Press.
- Ahmed, K. N. and Islam, W. (1988): A new record of the parasite *Rhabdepyris zeae* Waterston (Hymenoptera: Bethylidae) from Bangladesh and some aspects of its biology. Bangladesh Journal of Zoology, 16 (2):137-141.
- Aisagbonhi, C. I. (1988): Pest incidence in the marketed date palm fruits in Dutse, Kano state, Nigeria. Date palm J., 6 (1): 287-298.
- Aitken, A. D. (1963): A key to the larvae of some species of *Phycitinae* (Lepidoptera, Pyralidae) associated with stored products, and of some related species. Bulletin of Entomological Research, 54:175-188.
- Aitken, A. D. (1975): Insect Travellers. Volume 1. Coleoptera. Technical Bulletin 31, HMSO, xvi, Ministry of Agriculture Fisheries and Food: London, UK.
- Al-Abbasi, S. H. (1988): Antennal sense organs of *Ommatissus binotatus* De Berg (Tropiduchidae: Homoptera). Date Palm J. 6 (2): 426-431.
- Al-Abdulmohsin, A. M. (1987): First record of Red date palm weevil in Saudi Arabia. Arab World Agriculture 3: 15-16.

References

- Al-Azawi, A. F. (1986): A survey of insect pests of date palm in Qatar. Date Palm Journal (4): 247-266.
- Alexopoulou, V. P.; Wourikis, A. P. and Bouchelos, Th. K. (1986): *Coccotrypes dactyliperda* Fabr., a new species in the Greek fauna. New records. Chronika Benaki phytopathological Institute, 15 (1): 91-93.
- Alfieri, A. (1934): Sur une nouvelle maladie du dattier. Bulletin de la Société Entomologique d'Egypte 18, 445-448.
- Alfieri, M. A. (1976): The Coleoptera of Egypt. Soc. Ent. Egypt, (5): 361 pp.
- Al-Hafidh, E. M. T. (1979): Studies on some date insects. M. Sc. Thesis, Alexandria Univ., Egypt.
- Al-Hafidh, E. M. T. (1985): The integrated of *Nosema whitei* and some insecticides on *Tribolium castaneum*. Ph. D. Thesis, Newcastle upon Tyne Univ., UK.
- Al-Hafidh, E. M. T.; Siwer, E. A. and Abdul-Ahad, E. (1975): Biological and ecological studies on date palm stem borer, *Jebusea hammerschmidti* Reich, in Iraq. The third international conference of date palm and dates, Baghdad, Iraq. (In Arabic).
- Al-Kaabi, M. (1993): New Pest Attacking Date Palm Trees in Mahdha. Internal Report to the Director of Agriculture and Animal Wealth in Buraimi. 17 August 1993, (in Arabic).
- Al-Khatiri, S. A. (2004): Report of integrated management of red palm weevil (*Rhynchophorus ferrugineus* F.) in Oman. In Arabic, 5 pp.
- Al-Khatiri, S. A. and Abdallah, F. F. (2003): Seasonal fluctuation of *Rhynchophorus ferrugineus* Olivier (Coleoptera: Curculionidae) in the Sultanate of Oman. International Scientific Meeting of the Date Palm, 19-22 Ragab 1424H-16-19 September 2003. College of Agriculture and Veterinary Medicine, King Saud University-Qassim, Ministry of High Education, Kingdom of Saudi Arabia, (1): 371-385.
- Amitava Konar (1998): Seasonal incidence of mealybug species on mandarin orange in Darjeeling, West Bengal. Horticultural Journal, 11 (2):105-110.
- Anitha, B.; Arivalagan, M.; Sundari, M. S. N. and Durairaj, G. (1999): Effect of alkaloid abrine, isolated from Abrus precatorius Linn. seeds on mealy bug, *Maconellicoccus hirsutus* Green. Indian Journal of Experimental Biology, 37 (4): 415-417.
- Anonymous (1992a): List of Agricultural Insects and Mites in Oman. Agricultural Research Annual Report, Directorate General of Agricultural Research, Rumais, Sultanate of Oman, 171-179.
- Anonymous (1992b): Studies of the susceptibility of Date cultivars to the old world mite *Oligonychus afrasiaticus*. Agricultural Research Annual Report, Directorate General of Agricultural Research, Rumais, Sultanate of Oman, 180-181.
- Anonymous (1992c): Evaluation of the susceptibility of Omani Date cultivars to Graphiola leaf spot. Agricultural Research Annual Report, Directorate General of Agricultural Research, Rumais, Sultanate of Oman, 134-136.
- Anonymous (1993): Status of Pests and Diseases in Oman, Plant Diseases in the Batinah. Agricultural Research Center (ARC), Rumais, Sultanate of Oman, No. (6 / 93 / 22), 150 pp.
- Anonymous (1994): Biological control of some important insect pests. Agricultural Research Annual Report, Directorate General of Agricultural Research, Rumais, Sultanate of Oman, 325-341.

References

- Anonymous (1997): Dates quality improvement in the Sultanate of Oman, a comprehensive document. Sultane Qaboos University, College of Agriculture, 81 pp.
- Arai, T. (1996): Temperature-dependent development rate of three mealybug species, *Pseudococcus citriculus* Green, *Planococcus citri* (Risso), and *Planococcus kraunhiae* (Kuwana) (Homoptera: Pseudococcidae) on Citrus. Japanese Journal of Applied Entomology and Zoology, 40 (1): 25-34.
- Arar, A. (1975): Soils, irrigation and drainage of the date palm. 3rd FAO Tech. Conf. on Imp. Date Produc., Proc. And Marktg. Paper No. A3.
- Archer, M. E. (1998): Taxonomy, distribution and nesting biology of *Vespa orientalis* L. (Hym., Vespidae). Entomologist's Monthly Magazine 134: 45-51.
- Asche, M. and Wilson, M. R. (1989): The palm-feeding planthopper genus *Ommatissus* (Homoptera: Fulgoroidea: Tropiduchidae). Systematic Entomology, (14): 127-147.
- Ashworth, J. R. (1993): The biology of *Ephestia elutella*. Journal of Stored Products Research, 29 (3): 199-205.
- Atif, M. E. and Nazeef, M. H. (1998): Date Palm, cultivated and its care and production in the Arab world. Monsha'a El-Maaref, Alexandria, Egypt, 756 pp., (In Arabic).
- Awadallah, A. M. (1966): The biology and control of the pomegranate fruitfly *Viracola livia* Klug, (Lepidoptera, Lycaenidae) Ph. D. thesis, Cairo University.
- Awadallah, K. T.; Tawfik, M. F. S. and El-Husseini, M. M. (1986a): Bio-cycle of the anthocorid predator *Xylocoris flavipes* (Reuter) in association with rearing on major pests of stored drug materials. Bulletin de la Société Entomologique d'Egypte, No. (66): 27-33.
- Awadallah, K. T.; Tawfik, M. F. S. and El-Husseini, M. M. (1986b): Effect of larval preys on the biocycle of *Xylocoris sordidus* (Reuter) (Anthocoridae, Hemiptera). Archiv für Phytopathologie und Pflanzenschutz, 22 (3): 237-242.
- Awadallah, K. T.; Tawfik, M. F. S.; Abou-Zeid, N. and El-Husseini, M. M. (1980): The life-history of *Dufouriellus ater* (Duf.) (Hemiptera: Anthocoridae). Bulletin de la Société Entomologique d'Egypte, No. (63): 191-197.
- Ayers, R. S. and Westcot, D. W. (1985): FAO Irrigation and Drainage Paper 29 Rev. 1, Water Quality for Agriculture.
- Azam, K. M. and Razvi, S. A. (2001): Infestation of red palm weevil, *Rhynchophorus ferrugineus* in relation to offshoots on date palm trunk and its management. Paper presented in the meeting on the Red Palm Weevil, *Rhynchophorus ferrugineus*, Muscat, Oman, 19-20 February, 2000.
- Bai, B.; Cobanglu, S. and Smith, S. M. (1995): Assessment of *Trichogramma* species for biological control of forest lepidopteran defoliators. Entomologia Experimentalis et Applicata 75, 135-143.
- Baker, E. W. and D. M. Tuttle, 1972. New species and further notes on the Tetranychidae mostly from the southwestern United States (Acarina: Tetranychidae and Tenuipalpidae). Smithsonian Contributions to Zoology, 116: 1-37.
- Baker, E.W. (1948): A new trichadenid mite which further indicates a phylogenetic relationship between the

References

- Tenuipalpidae and the Eriophyidae (Acarina). Proc. Ent. Soc. Wash., (50): 59-60.
- Banks, H. J. (1977): Distribution and establishment of *Trogoderma granarium* Everts (Coleoptera: Dermestidae); climatic and other influences. Journal of Stored Products Research, 13 (4): 183-202.
- Barker, P. S. (1968): Note on the bionomics of *Haemogamasus pontiger* (Berlese) (Acarina: Mesostigmata), a predator on *Glycyphagus domesticus* (DeGeer). Manitoba Entomologist, (2): 85-87.
- Barranco, P.; de la Pena, J. A.; Martin, M. M. and Cabello, T. (2000): Rango de hospedantes de *Rhynchophorus ferrugineus* (Olivier, 1790) y diámetro de la palmera hospedante. (Coleoptera: Curculionidae). Boletín de Sanidad Vegetal Plagas (26): 73-78.
- Bartlett B. R. (1978): Pseudococcidae. In: Clausen CP, ed. Introduced Parasites and Predators of Arthropod Pests and Weeds: a World Review. Agriculture Handbook No. 480, 137-170.
- Batra, R. C. (1972): Insect pests of date palm at Abohar and their control. Punjab Hort. J. (12): 44-45.
- Bedford, G. O. (1976): Use of a virus against the coconut palm rhinoceros beetle in Fiji. PANS (22): 11-25.
- Bedford, G. O. (1980): Biology, ecology and control of palm rhinoceros beetles. Annual Review of Entomology, (25): 309-339.
- Bell, C. H. (1975): Effects of temperature and humidity on development of four pyralid moth pests of stored products. Journal of Stored Products Research, 11(3/4): 167-175.
- Bénassy, C. (1990): Armoured scale insects, their biology, natural enemies and control. Vol. 4B. World Crop Pests. Elsevier, Amsterdam, the Netherlands: 585-591.
- Bénassy, C.; Deportes, L.; Onillon, J. C. and Panis, A. (1976): Orientation towards integrated control in citrus culture in south-east France. Pepinieristes Horticulteurs Maraîchers, 167: 41-48.
- Benbrook, C. (2004): Executive Summary, Genetically Engineered Crops and Pesticide Use in the United States: The First Nine Years. October 25. www.biotech-info.net
- Bernays, E. A. and Chapman, R. F. (1978): Plant chemistry and acridoid feeding behaviour. In: Harborne, ed. Biochemical aspects of plant and animal co-evolution. Symposium of the Phytochemical Society of Europe. London, UK: Centre for Overseas Pest Research, (5): 99-151.
- Besri, M. (1983): Importance of Bayoud Disease (*Fusarium oxysporum* f.sp. *alpedenis malencon*) of Date Palm in Morocco. The First Symposium on The Date Palm in Saudi Arabia, March, 23-25, 1982, 424-430.
- Betrem, J. G. (1936): Gegevens omtrent de biologie van de dompololanluis en de lamtorolois', arch. Koffiecult. Ned. Indie, (10): 43-84.
- Bhatia, S. K. and Nadarajan, L. (1978): Inheritance of black body colour in a mutant of *Tribolium castaneum* (Herbst) with notes on its biology. Journal of Entomological Research, 2 (2): 219-221.
- Bishop, G. W. (1960): Taxonomic observations on the larvae of the three American *Cryptolestes* (Coleoptera: Cucujidae) that infest stored grain. Annals of the Entomological Society of America, (53): 8-11.

References

- Bitton, S. and Nakache, Y. (2000): Red palm weevil in Israel. Proceeding Date Palm International Symposium February 22-25, 2000, Republic of Namibia.
- Blancaver, R. C.; Abad, R.G.; Pacumbaba, E. P. and Mordeno, J. C. (1977): Guidebook on Coconut Pests and Diseases. Philippine Coconut Authority, Davao Research Center - Philippine.
- Bliss, D. E. (1936): Rhizosis, a recently discovered disease date palm. Date Grower's Inst Rpt. (13): 4-6.
- Bliss, D. E. (1937): Crosscuts in the Fruitstalks of date plant. Ann. Date Growers, Instit. (14): 8-11.
- Bliss, D. E. (1944): Omphalia root rot of the date palm. Hilgardia, (16): 15-124.
- Blumberg, D. and Kehat, M. (1982): Biological studies of the date stone beetle, *Coccotrypes dactyliperda*. Phytoparasitica (10): 73-78.
- Boake, C. R. B. and Wade, M. J. (1984): Populations of the red flour beetle *Tribolium castaneum* (Coleoptera: Tenebrionidae) differ in their sensitivity to aggregation pheromones. Environmental Entomology, 13 (5): 1182-1185.
- Bodenheimer, F. S. (1937): Prodromus Faunae Palastinae. Meu. Inst. Egypt, 33 pp.
- Booth, R. G. and Pope, R. D. (1986): A review of the genus *Cryptolaemus* (Coleoptera: Coccinellidae) with particular reference to the species resembling *C. montrouzieri* Mulsant. Bulletin Entomological Research (76): 701-717.
- Booth, R. G.; Cox, M. L. and Madge, R. B. (1990): IIE Guides to Insects of Importance to Man. 3. Coleoptera. CAB International, Wallingford (GB).
- Boyden B. L. (1941): Eradication of the Parlatoria date scale in the United States. USDA, Miscellaneous Publication, 433.
- Bright, D. E. (1981): Eye reduction in a cavernicolous population of *Coccotrypes dactyliperda* Fabr. (Coleoptera: Scolytidae). Bull. Coleopterists, 35 (1): 117-120.
- Brun, J. and Laville, E. (1965): Observations sur un dépérissement de la couronne foliaire et du bourgeon terminal du palmier dattier en République Islamique de Mauritanie. Fruits (20): 391-397.
- Bult, J.; Louvet, J.; Bouhot, D. and Toutain, G. (1967): Recherches sur les fusarioses. I. Travaux sur le bayoud Fusariose vasculaire du palmier dattier en Afrique du Nord. Ann. Epiphyt. 18 (2): 213-239.
- CAB Abstracts (1973-1998): Data mined from CAB Abstracts database, years 1973 to 1998. Wallingford, UK: CAB International.
- CABI (CAB International). (2005): Crop protection compendium CD-ROM. Wallingford, Oxon, U.K.
- Calcat, A. (1959): Disease and pests of date palm in the Sahara and North Africa. FAO plant Protect. Bull. (8): 5-10.
- Carpenter, J. B. (1975): Notes on date culture in the Arab Republic of Egypt and People's Republic of Yemen. Date Growers' Inst. Report (52): 18-24.
- Carpenter, J. B. and Elmer, H. S. (1978): Pests and Diseases of the Date Palm. Agriculture Handbook No. 527, US Department of Agriculture, Washington, DC.

References

- Carpenter, J. M. and Kojima, J. (1997): Checklist of the species in the subfamily Vespinae (Insecta: Hymenoptera: Vespidae). Natural History Bulletin of Ibaraki University (1): 51-92.
- Chase, A. R. and Broschat, T. K. (1991): Diseases and Disorder of Ornamental Palms. The American Phytopathological Society, USA. 56 pp.
- Chazeau, J., 1974a. Développement et fécondité de *Stethorus madecassus* Chazeau (Coléoptères, Coccinellidae) élevé en conditions extérieures dans le sud-ouest de Madagascar. Cahiers ORSTOM, Série Biologie (25): 27-33.
- Chazeau, J., 1974b. Evaluation de l'action prédatrice de *Stethorus madecassus* (Coléoptères, Coccinellidae) sur *Tetranychus neocaledonicus* (Acarien Tetranychidae). Entomophaga (19): 183-193.
- CIE (1967): Distribution Maps of Plant Pests, No. 54. Wallingford, UK: CAB International.
- Coudin, B. and Galvez, F. (1976): Biology of the date palm mite *Oligonychus afrasiaticus* (Mc Gregor) in Mauritania. Fruits (France) 1976, 31 (9): 543-550.
- Cox, J. M. (1989): The mealybug genus *Planococcus* (Homoptera: Pseudococcidae). Bulletin of the British Museum (Natural History), Entomology, 58 (1): 1-78.
- Cox, M. L. (1993): Red palm weevil *Rhynchophorus ferrugineus* F. in Egypt, FAO Plant Protection Bulletin (41): 30-31.
- Daniel, M. (1981): Bionomics of the predaceous mite *Amblyseius channabasavanni* (Acari: Phytoseiidae), predaceous on the palm mite *Raoiella indica*. In: 1st Indian Symposium in Acarology (G. P. Channabasvanna, ed.). April 23-25, Bangalore, India, 167-173.
- Darley, E. F. (1964): Second report on the bending of tops of Barhee date Palms. Date Grower's Inst. Rpt., 41-61.
- Darley, E. F. and Wilbur, W. D. (1955): Results of experiments on control of fruit spoilage of Deglet Noor and Saidy dates in California, 1935-1954. Ann. Date Grower's Instit. (32): 14-15.
- Darley, E. F.; Nixon, R. W. and Wilbur, W. D. (1960): An unusual disorder of Barhee date palms. Ann. Date Grower's Instit., (37): 10-12.
- Davidson, J. A. and D. R. Miller (1990): Ornamental plants. In: D. Rosen (ed.), Armoured scale insects, their biology, natural enemies and control. Vol. 4B. Elsevier, Amsterdam, the Netherlands: 603-632.
- De Bergevin, E. (1930): Note sur trois espèces d'hémiptères recueillis en Egypte et description d'une nouvelle espèce d'*Urentius* (hémiptère Tingitidae) et d'une nouvelle variété *d'Ommatissus binotatus* Fieb. (homoptère Cixiidae). Bulletin de la Société Entomologique d'Egypte. Le Caire: 17-20.
- De Boer, A. J. and Duffels, J. P. (1996): Historical biogeography of the cicadas of Wallacea, New Guinea and the West Pacific: a geotectonic explanation. Paleogeography, Palaeoclimatology, Palaeoecology 124: 153-177.
- Dechambre R. P. (1983): Contribution to the phylogenetic study of the *Oryctes* of the Madagascan region; use of the propygidal stridulatory stria (Col. Dynastidae). Bulletin de Société Entomologique de France, 88(5/6): 436-448.
- Dembilio, Ó.; Jacas, J. A. and Llácer , E. (2009): Are the palms *Washingtonia filifera* and *Chamaerops*

References

- humilis* suitable hosts for the red palm weevil, *Rhynchophorus ferrugineus* (Col. Curculionidae). Journal of Applied Entomology. Vol. 133, Issue (7): 565-567.
- Dick, G. L.; Buschman, L. L. and Ramoska, W. A. (1992): Description of a species of *Neozygites* infecting *Oligonychus pratensis* in the western great plains of the United States. Mycologia, (84): 729-738.
- Djerbi, M. (1980): Report on duty travel to Tunis, Baghdad and Pakistan. FAO Regional Project for Palm and Dates Research Center in the Near East and North Africa Baghdad, Iraq, 10 pp.
- Djerbi, M. (1983): Diseases of the Date Palm. FAO Regional Project for Palm and Dates Research Center in the Near East and North Africa, 106 pp.
- Djerbi, M.; Aouad, L.; Filali, H.; Saaidi, M.; Chtioui, A.; Sedra, M. H.; Allaoui, M.; Hamdaoui, T. and Oubrich, M. (1986): Preliminary results of selection of high quality Bayoud resistant clones among natural date palm population in Morocco. The second symposium on the date palm, Saudi Arabia, March 3-6. Vol. (11): 383-399.
- Doumandji, M. B. (1977): Les pyralis des dates stokees. Extracted from Ann. De l'Inst. National Agronom. (El-Harrach), Algeria, 7 (1): 1-58.
- Dowson, V. H. W. (1936): A serious pest of date palms. *Ommatissus binotatus* Fieb. (Homoptera: Tropiduchidae). Tropical Agriculture (Trinidad) (13): 180-191.
- Dowson, V. H. W. and Pansiot, F. P. (1965): Improvement of date palm growing. FAO, Agri. Dev. Paper No. 72.
- Ebrahim, H. M. (2000): Large-scale mass production of the predatory mite, *Phytoseiulus persimilis* A. H. and the predatory insect, *Stethorus gilvifrons* (Mulsant) reared on *Tetranychus urticae* Koch in greenhouses. Journal of the Egyptian German Society of Zoology, HHP: www.mideastnet.com / egygersoc, Email: (egygersoc@arabmarket.com), Vol. 32 (E), Entomology, 51-58, March 2000.
- El Ezaby, F. (1997): A biological in vitro study on the red Indian date palm weevil. Arab Journal of Plant Protection, 15 (2): 84-87.
- Elarosi, H.; El-Said, H.; Nagieb, M. A. and Jaheen, N. (1983): Al-wijam, Decline date palm Disease. The First Symposium on the Date Palm in Saudi Arabia, March: 23-25, 1982, 388-403.
- El-Baker, A. J. (1952): Report to the Govt. of Saudi Arabia on date cultivation in Saudi Arabia. Rome FAO/ EPTA Report No. 31.
- Elder, R. J. and Smith, D. (1995): Mass rearing of *Aonidiella orientalis* (Newstead) (Hemiptera: Diaspididae) on butternut gramma. Journal of the Australian Entomological Society, 34 (3): 253-254.
- Elder, R. J.; Smith, D. and Bell, K. L. (1998): Successful parasitoid control of *Aonidiella orientalis* (Newstead) (Hemiptera: Diaspididae) on Carica papaya L. Australian Journal of Entomology, 37 (1): 74-79.
- El-Haidari, H. S. (1979): Report to the governments of Kuwait, Bahrain, Qatar and UAE. Regional project for palm and dates Res. Cent. Baghdad, Iraq.
- El-Haidari, H. S. (1980): Palm and Date Arthropod Pests in the Near East and North Africa. Regional project for palm and dates research center in the Near East and North Africa. Baghdad, Iraq.
- El-Haidari, H. S. (1982): New records of Dubas bug (*Ommatissus binitatus lybicus* de Bergevin) on date palms in Sudan. Date Palm Journal, 1 (2): 308.

References

- El-Haidari, H. S. and Al-Hafidh, E. M. T. (1986): Palm and Date Arthropod Pests in the Near East and North Africa. Regional project for palm and dates research center in the Near East and North Africa. Al-Watan Press, Lebanon. 126 pp.
- El-Haidari, H. S. and Al-Tigany, M. A. (1977): Studies on the control of Dubas bug on date palm and Citrus Black Fly on citrus in Sultanate of Oman. Arab Organization for Agriculture Development, Khartoum, Sudan.
- El-Haidari, H. S.; Al-Safi, G.; Michael, N. and Sallam, N. (1982): Effect of constant temperatures on the development of the dust date palm mite *Oligonychus afrasiaticus*. Annual Book Plant Protection Research 2 (1): 165-167.
- El-Haidari, H. S.; Al-Zidjali, T. S. and Al-Bussaidi, A. A. (1992): Evaluation of the susceptibility of Omani Datepalm to Lesser Date Moth *Batrachedra amydraula* infestation. Agric. Res. Ann. Rep., Dept. Agric. Res., Rumais, Oman, 182-183.
- El-Haidari, H. S.; Thiab, E. M.; Hussain, I. A. and Abdul Wahab, W. A. (1975): Studies on the biology of lesser date moth. *B. amydraula* in Iraq. The Third International Palm and Dates Conference, Baghdad, Iraq.
- El-Kareim, A. I. A. (1998): Swarming activity of the adult males of *Parlatoria* date scale in response to sex pheromone extracts and sticky colour traps. Archives of Phytopathology and Plant Protection 31, 301-307.
- Elliot, M. L. (2006): Thielaviopsis trunk rot of palm. Fact Sheet pp. 219, Plant Pathology Department, Florida Cooperative Extention Service, Institute of Food and Agricultural Sciences, University of Florida.
- Ellis, P. E. and Ashall, C. (1957): Field studies on diurnal behaviour, movement and aggregation in the Desert Locust, *Schistocerca gregaria* Forskal. Anti-Locust Research Centre: London Bulletin No. (25): 94 pp.
- El-Naggar, M. Z.; Etman, A. A. M. and El-Hemeasy, A. H. (1985): Distribution of the desert subterranean termit, *Amithermes desertorum* (Deseneux) attacking date palm trees (*Phoenix dactylifera* L.) in Aswan. Bull Fac. Agric, Cairo Univ. 36 (1): 585-592.
- El-Sayed, S. A. and Baeshin, N. A. (1983): Feasibility of disinfestations of date fruits produced in Saudi Arabia by Gamma irradiation. The First Symposium on the Date Palm in Saudi Arabia, March: 23-25, 1982, 342-350.
- El-Sufti, R.; Al-Awash, S. A.; Al-Amiri, A. M.; Shahdad, A. S.; Al-Bathra, A. H.; and Musa, S. A. (2007): Biological control of red palm weevil, *Rhynchophorus ferrugineus* (Col.: Curculionidae) by the entomopathogenic fungus *Beauveria bassiana* in United Arab Emirates. Acta Horticulturae 736: 399-404.
- Elwan, A. A. (2000): Survey of the insect and mite pests associated with date palm trees in Al-Dakhliya region, Sultanate of Oman, Egypt. J. Agric. Res. 78 (2): 653-664.
- Entwistle, P. F. (1972): Pests of coffee. London, UK: Longman Group Limited.
- EPPO (2006): PQR database (version 4.5). Paris, France: European and Mediterranean Plant Protection Organization. (www.eppo.org).
- EPPO (2008): Data sheet on quarantine pests. *Rhynchophorus ferrugineus*. Eppo (European and Mediterranean Plant Protection Organization) Bull., (38): 55-59.

References

- Eriksen, J. and Sargeant, D. E. (2000): Oman Bird List. Edition 5, Oman Bird Records Committee, P.O. Box 246, Muscat 113, Sultanate of Oman, 168 pp.
- Etienne, J. and Fletchmann, C. H. W. (2006): First record of *Raoiella indica* (Hirst, 1924) (Acari: Tenuipalpidae) in Guadeloupe and Saint Martin, West Indies. Internat. J. Acarol. (32): 331-332.
- FAO (1978): Research on the control of the coconut palm rhinoceros beetle, phase II. Fiji, Tonga and Western Samoa. Tech. Res., Rome, 94 pp.
- FAO (1982): Date palm Date palm production and protection. FAO plant production and protection paper, No. 35, Food and Agriculture Organization of the United Nation, Rome, 294 pp.
- FAO (1993): Date palm Date palm products. FAO Agricultural Services Bulletin, No. 101, Food and Agriculture Organization of the United Nation, Rome, 216 pp.
- FAO (1995): Report of the Expert Consultation on Date Palm – Pest Problems and their control in the Near East. (22-26 April, 1995); Al-Ain, United Arab Emirates.
- FAO (1999): Date palm cultivation. FAO Plant Production and Protection Paper. No. 156, Food and Agriculture Organization of the United Nation, Rome, 287 pp.
- Farid, A. (1994): Study on bio-ecology and control of *Aonidiella orientalis* in Jiroft and Hormozgan. Applied Entomology and Phytopathology, 61(1&2): 29, Ar 96-105.
- Farr, D. F.; Bills, G. F.; Chamuris, G. P. and Rossman, A.Y. (1995): Fungi on plants and plant products in the United States. American Phytopathological Soc. St. Paul MN.
- Fawcett H. S. and Klotz, L. J. (1932): Diseases of the date palm, *Phoenix dactylifera* L. Calif. Agric. Exp. Sta. Bull. 522, 47 pp.
- Fawcett, H. S. (1930): An offshoot and leafstalk disease of Date palms due to *Diplodia*. Phytopathology, 1930 Vol. 20, No. (4): 339-344.
- Feather, T. V.; Ohr, H. D. and Munnecke, D. E. (1979): Wilt and dieback of canary island palm in California. California Agriculture (33):19-20.
- Feather, T. V.; Ohr, H. D.; Munnecke, D. E. and Carpenter, J. B. (1989): The occurrence of *Fusarium oxysporum* on *Phoenix canariensis*, a potential danger to date production in California.
- Ferron, P. (1985): Comperhensive Insect Physiology, Biochemistry, Pharmacology, Vol. 12, Insect Control, Pergamon Press, 849 pp.
- Fiaboe, K. K. M.; Gondim, M. G. C.; De Moraes, G. J.; Ogol, C. K. P. O. and Knapp, M. (2007): Bionomics of the acarophagous ladybird beetle *Stethorus tridens* fed *Tetranychus evansi*. Journal of Applied Entomology 131, 355-361.
- Fieber, F. X. (1875): Les Cicadines d'Europe d'après les originaux et les publications les plus récentes. Première partie. Revue et Magasin de Zoologie pure et appliquée. Paris. (Ser. 3) (3): 288-416.
- Figueroa, W. (1990): Biocontrol of the banana root borer weevil, *Cosmopolites sordidus* with steiner nematid nematodes. Journal of Agriculture, University of Puerto Rico, (74): 15-19.
- Flinn, P. W. and Hagstrum, D. W. (1995): Simulation model of *Cephalonomia waterstoni* (Hymenoptera: Bethylidae) parasitizing the rusty grain beetle (Coleoptera: Cucujidae). Environmental Entomology, 24 (6):1608-1615.

References

- Flint, M. L. and Dreistadt, S. H. (1998): Natural enemies handbook. The illustrated guide to biological pest control. University of California, Statewide Integrated Pest Management Project. Uc Division of Agriculture and Natural Resources, and the University of California Press, 154 pp.
- Fontes, L. S.; Arthur, V.; Wiendl, T. A. and Wiendl, F. M. (1996): Gamma radiation effects on pupae of *Tribolium castaneum* (Herbst). Journal of Nuclear Agriculture and Biology, 25 (3):176-179.
- Forst, S. and Clarke, D. (2002): Bacteria-Nematodes symbiosis. In Gaugler, R. entomopathogenic nematodes. Chapter 3. CAB International, United Kingdom.
- Forst, S.; Dowds, B.; Boemare, N.; and Stackebrandt, E. (1997): *Xenorhabdus* spp. And *Photorhabdus* spp. that kill bugs. Annual Review of Microbiology. (51): 47-72.
- Friedrichs, K. (1913): Ueber der Bekämpfung des Nashornkafers (*Oryctes rhinoceros*) in Samoa. Tropenpflanzer, 17 (10): 538.
- Fronteddu, F.; Canu, D.; D'Amico, R.; Delpiano, N.; Fancello, F. and Nanni, G. (1996): Applications of integrated control methods in citrus cultivation: biotechnical control against *Ceratitis capitata* and biological control of *Planococcus citri*. Informatore Fitopatologico, 46 (11): 34-39.
- Funasaki, G. Y.; Nakahara, L. M. and Kumashiro, B. R. (1988): Introductions for biological control in Hawaii: 1985 and 1986. Proceedings of the Hawaiian Entomological Society, No. (28): 101-104.
- Gallagher, M, and Woodcock, M. W. (1980): The Birds of Oman. Quarter Book, London, UK.
- Gassouma, S. M. (2005): Pests of Date Palm, *Phoenix dactylifera*, (www.uae.gov.ae/uaeagricent/palmtree2/chap8.stm).
- Gaugler, R. and Kaya, H. K. (1990): Entomopathogenic Nematodes in Biological Control. Boca Raton, FL: CRC Press.
- Geijskes, D. C. (1939): Beitrage zur kenntnis der europäischen Spinnmilben (Acari: Tetranychidae) mit besonderer Berücksichtigung der niederlandischens Arten. Meded. Landbouwhooge School. Wageningen. 42 (2): 1-68.
- Gerson, U.; Venezian, A. and Blumberg, D. (1983): Phytophagous mites on date palms in Israel. Fruits (38): 133-135.
- Ghabbour, M. W.; Mohammad, Z. K. and Elwan, E. A. (1996): *Fiorinia linderae* Takagi new record in Oman (Homoptera: Diaspididae). J. Egyptian German Soc. Zoology Vol. 19 (E), Entomology, 51-58.
- Gharib, A. (1968): *Batrachedra amydraula* Meyr. (Super-Family: Gelechioidea, Family: Momphidae (Cosmopterygidae)). Entomol. Phytopath. Appl., (27): 63-66.
- Gharib, A. (1969): *Arenipses sabella* Hamp. (Lepidoptera, Pyralidae). Entomol. Phytopath. Appl., (28): 34-35.
- Gharib, A. (1973): *Parlatoria blanchardi* Targ. (Homoptera-Diaspididae). Entomologie et Phytopathologie Appliquées (34): 10-17.
- Gharib, A. (1976): *Paratetranychus (Oligonychus) afrasiaticus* (McG.) Tetranychidae Entomol. Phytopath. Appl., 26: 27-30.
- Ghauri, M. (1962): The morphology and taxonomy of male scale insects (Homoptera, Coccoidea). British Museum (Natural History). Adlard and Son, Dorking, UK, 221 pp.

References

- Ghose, S. K. (1971): Morphology of various instars of both sexes of the mealybug, *Maconellicoccus hirsutus* (Green) (Pseudococcidae: Hemiptera). Indian Journal of Agricultural Sciences, 41 (7): 602-611.
- Ghose, S. K. (1972): Biology of the mealybug, *Maconellicoccus hirsutus* (Green) (Pseudococcidae, Hemiptera). Indian Agriculturist, (16): 323-332.
- Gill, R. J. (1997): The scale insects of California. Part 3. The armoured scales (Homoptera: Diaspididae). Technical Series in Agricultural Biosystematics and Plant Pathology, No 3. Sacramento, USA: Department of Food and Agriculture.
- Gillett, S. D. (1973): Social determinants of aggregation behaviour in adults of the desert locust. Anim. Behav. (21): 599-606.
- Glazer, I.; Salame, L.; Goldenberg, S. and Blumberg, D. (1999): Susceptibility of sap beetles (Coleoptera: Nitidulidae) to entomopathogenic nematodes. Biocontrol Science and Technology, 9 (2): 259-266.
- Glover, P. M. (1933): *Aspidiotus (Furcaspis) orientalis* Newstead, its economic importance in lac cultivation and its control. Indian Lac Research Institute Bulletin, (16): 1-22.
- Golemanski, V. G. and Dukhlinska, D. D. (1982): Unicellular parasites of insect pests in Bulgaria. I. Composition and distribution of Sporozoa and Microsporidia in grain storehouse insect pests. Acta Zoologica Bulgarica, (20): 26-37.
- Gorick, B. D. (1980): Release and establishment of the baculovirus disease of *Oryctes rhinoceros* (L.) (Coleoptera: Scarabaeidae) in Papua New Guinea. Bull. Ent. Res. (70): 445-453.
- Gough, L. H. (1913): Entomological notes. Agric. Jour. Of Egypt, Cairo, Vol. 3 No. (2): 103-106.
- Gravitz, N. and Wilson, C. (1968): A sex pheromone from the citrus mealybug. Journal of Economic Entomology, (61): 1458-1459.
- Greathead, D. J. (1977): Report on a visit to the Sultanate of Oman. Commonwealth Institute of Biological Control.
- Greathead, D. J. and Greathead, A. H. (1992): Biological control of insect pests by insect parasitoids and predators: the BIOCAT database. Biocontrol News and Information, 13 (4): 61N-68N.
- Gries, G.; Gries, R.; Perez, A. L.; Gonzalez, L. M.; Pierce, H. D.; Oehlschlager, A. C.; Rhainds, M.; Zebeyou, M. and Kouame, B. (1994): Ethyl propionate, synergistic kairomone for African palm weevil, *Rhynchophorus phoenicis* L. (Coleoptera: Curculionidae). Journal of Chemical Ecology, (20): 889-897.
- Gullan, P. J. (2000): Identification of the immature instars of mealybugs (Hemiptera: Pseudococcidae) found on citrus in Australia. Australian Journal of Entomology (2000), (39): 160-166.
- Gunawardena, N. E. and Bandarage, U. K. (1995): 4-Methyl-5-nonal (ferrugineol) as an aggregation pheromone of the coconut pest, *Rhynchophorus ferrugineus* F. (Coleoptera: Curculionidae): synthesis and use in a preliminary field assay. Journal of the National Science Council of Sri Lanka (23): 71-79.
- Gupta, Y. N. (2001): A conspectus of natural enemies of phytophagous mites and mites as potential biocontrol agents of agricultural pests in India. In: International Congress of Acarology, R. Halliday, D. Walter, H. Proctor, R. Norton and M. Colloff, eds. 10. Collingwood, Australia. CSIRO Publishing, 484-497.

References

- Haines, C.P. (1981): Insects and arachnids from stored products: a report on specimens received by the Tropical Stored Products Centre 1973-77. Tropical Products Institute Report L54, September 1981, 73 pp.
- Haines, C.P. (1991): Insects and Arachnids of Tropical Stored Products: Their Biology and Identification (A Training Manual), 2nd edition. Chatham, UK: Natural Resources Institute.
- Hall, A.V. (1988): Pest control in herbaria. *Taxon*, 37 (4): 885-907.
- Hall, W. J. (1921): The hibiscus mealybug (*Phenacoccus hirsutus* Green). Bulletin Ministry of Agriculture Egypt Technical and Scientific Service Entomological Section, (17): 1-28.
- Hallett, R. H.; Oehlschlager, A. C.; Gries, G.; Angerilli, N. P. D.; Shareqi, R. K.; Gasouma, M. S. and Borden, J. H. (1993): Field testing of aggregation pheromones of two Asian palm weevils. In: Proceedings of the International Oil Palm Congress. Palm Oil Research Institute of Malaysia, Kuala Lumpur, Malaysia, 661-668.
- Hamdi, A. N. (1998): Report on present status of Red Palm Weevil and date palm borers in Bahrain. Arab Organisation for Agricultural Development (AOAD) Bulletin, 1-5 (in Arabic).
- Hammad, S. M.; Kadous, A. A. and Ramadan, M. M. (1981): Studies on insects and mites attacking date palm in the Eastern province of Saudi Arabia. Proceedings of Saudi Biological Society, (5): 251-268.
- Hammad, S. M.; Kadous, A. A. and Ramadan, M. M. (1983a): Seasonal abundance of the scale insect *Aonidiella orientalis* (Newstead) on Rizez variety of date palm in Al-Qatif region, Saudi Arabia. Proceedings of the First Symposium on the Date Palm, in Saudi Arabia, 23-25 March 1982, King Faisal Univ., Al-Hassa, Kingdom of Saudi Arabia, 298-305.
- Hammad, S. M.; Kadous, A. A. and Ramadan, M. M. (1983b): Predators and Parasites of date palm insects in Al-Hassa and Al-Qatif regions, Eastern province, Saudi Arabia. Proceedings of the First Symposium on the Date Palm, in Saudi Arabia, 23 – 25 March 1982, King Faisal Univ., Al-Hassa, Kingdom of Saudi Arabia: 322-341.
- Hammes, C. (1971): Multiplication et introduction d'un virus d'*Oryctes rhinoceros* à l'île Wallis. Comptes rendus de l'Academie des Sciences de Paris, Serie D, 273: 1048-1050.
- Hammouda A. M. (1985): Diseases of Coconut Palm and its Control Measures. Directorate General of Agriculture, Ministry of Agriculture & Fisheries, Sultanate of Oman, 52 pp. (In Arabic).
- Hammouda A. M.; Mekki, M. A. and El-Abri, A. S. (1998): Science of Fruit Orchards, Part Two, Date Palm, Vol. II, The Dates Varieties of Sultanate of Oman. Diwan of Royal Court, Directorate General of Agriculture & Veterinary, 642 pp. (In Arabic).
- Hanna, A. D. (1939): The pomegranate fruit butterfly *Viracolaliavia* Klug. Morphology, life history and control. Bull. Ministry of Agric. Egypt. No. 186, 54 pp.
- Hanounik, S. B. (1998): Steinemematis and heterorhabditids as biological control agent for the red palm weevil (*Rhynchophorus ferrugineus* Oliv.). Sultan Qaboos University, Journal for Scientific Research, Agricultural Sciences (3): 95-102.
- Hanounik, S. B.; Salem, M.; Hegazy, G.M.; Al-Muhana, O.; Al-Hagi, M. and Al Dhaher, H. (2000): Development of a new food baited aggregation pheromone/kairomone trapping system for the red

References

- palm weevil *Rhynchophorus ferrugineus* (Oliv.); Proceeding 1st International Workshop on control of Date Palm Red Weevil, King Faisal University, Saudi Arabia, 20-22, November (2000): 113-124.
- Haramoto, F. H. (1969): Biology and control of *Brevipalpus phoenicis* (Geijskes) (Acarina: Tenuipalpidae). Hawaii Agricultural Experiment Station Technical Bulletin No. (68): 1-63.
- Harrison, D. L. (1972): The Mammals of Arabia. 3. Lagomorpha. Rodentia. Benn. London, 382-670.
- Harrison, D. L. (1975): The Oman Flora and Fauna Survey, 1975. J. of Omani Studies I: 181-186.
- Hassan, B. R.; Al-Jaboory I. J.; Al-Rubeai, H. F. and Viggiani. G. (2004): *Pseudoligosita babylonica* n. sp. (Hymenoptera: Trichogrammatidae), egg parasitoid of *Ommatissus lybicus* Bergevin (Homoptera: Tropiduchidae) in Iraq. Bollettino del Laboratorio di Entomologia Agraria Filippo Silvestri (59): 75-78.
- Hegazi, E. M.; Agamy, E.; Hassan, S.; Hertz, A.; Khafagi, W.; Showel, S.; Abo-Abdala, L.; Ziton, A.; El-Said, S.; El-Shazly, A.; El-Menshawy, A.; Karam, H.; Khamis, N. and El-Kemny, S. (2004): Successful application of inundative releases of the egg parasitoid *Trichogramma evanescens* West. to control the olive moth, *Prays oleae* Bern. 1st Arab Conference of Applied Biological Pest Control, Cairo, Egypt, 5-7 April, 2004.
- Helle, W. and Sabelis, M. W. (1985a): Spider Mites: Their Biology, Natural Enemies, and Control, vols. 1 A. Elsevier, Amsterdam, 403 pp.
- Helle, W. and Sabelis, M. W. (1985b): Spider Mites: Their Biology, Natural Enemies, and Control, vols. 1 B. Elsevier, Amsterdam, 458 pp.
- Hilgeman, R. H. and Smith, J. G. (1937): The crude fat content of date skins correlated with moisture damage. Date Growers' Inst. Rpt., (14): 16-17.
- Hill, D. S. (1975): Agricultural insect pests of the tropics and their control. Cambridge University Press, 516 pp.
- Hill, D. S. (1983): In Agricultural Insect Pests of the Tropics and Their Control, 2nd Edition. Cambridge University Press, 746 pp.
- Hill, D. S. and Waller, J. M. (1982): Pests and diseases of tropical crops. Vol. 1. Principles and methods of control. London, UK: Longman.
- Hinton, H. E. (1945): A monograph of the beetles associated with stored products. Vol. 1. London, UK: British Museum (Natural History).
- Hoddle, M. S. (1998): Biology and management of the perseae mite. California Avocado Society Yearbook, (82): 75-85.
- Hodges, R. J.; Benton, F. P.; Hall, D. R. and Serôdio Rdos, S. (1984): Control of *Ephestia cautella* (Walker) (Lepidoptera: Phycitidae) by synthetic sex pheromones in the laboratory and store. Journal of Stored Products Research, 20 (4): 191-197.
- Hollis, J. P. and Johnston, T. (1957): Microbiological reduction of nematode populations in water-saturated soils. Phytopathology, 47:16.
- Howard, F. W.; Moore, D.; Giblin-Davis R. M. and Abad, R. G. (2001): Insects on Palms. CABI Publishing. CAB International, Wallingford, Oxon OX10 8DE, UK, 400 pp. Web site: www.cabi.org
- Howe, R.W. (1956): The biology of the two common storage species of *Oryzaephilus* (Coleoptera, Cucujidae). Annals of Applied Biology, 44 (2): 341-355.

References

- Hoy, M. A. (1984): Managing mites in almonds: an integrated approach. University of California, Davis, Integrated Pest Management Publication 1, 26 pp.
- Hüger, A. M. (1973): A virus disease of the Indian rhinoceros beetle, *Oryctes rhinoceros* (Linnaeus), caused by a new type of insect virus, *Rhabdionvirus oryctes* gen. N., sp. N. Journal of Invertebrate Pathology (8): 38-51.
- Hunter, C. D. (1997): Suppliers of beneficial organisms in north America. California Environmental Protection Agency, Department of Pesticide Regulation, Environmental Monitoring and Pest Management Branch, 1020 N Street, Sacramento, California 95814-5604. This booklet is electronically available, Natural Enemy Supplier Handbook. www.cdpr.ca.gov/docs/ipminov/bensupp.htm
- Hussain, A. A. (1963a): Biology and control of the dubas bug, *Ommatissus binotatus lybicus* de Berg. (Homoptera, Tropiduchidae) infesting date palms in Iraq. Bulletin of Entomological Research (53): 737-745.
- Hussain, A. A. (1963b): Notes on borers of date palms in Iraq. Bulletin of Entomological Research (54): 345-348.
- Hussain, A. A. (1967): Biology of *Tenuipalpus eriophyoides* Baker on date palms in Iraq. Bull. Soc. Ent. Egypt, (51): 31-32.
- Hussain, A. A. (1969): Biology of *Paratetranychus afrasiaticus* McGr. Infesting date palm in Iraq. Bull. Sci. Ent. Egypte, (53): 221-225.
- Hussain, A. A. (1974): Date Palms and Dates with their Pests in Iraq. 1st Ed., University of Baghdad, Ministry of Higher Education and Scientific Research, Baghdad, 164 pp.
- Hussain, A. A. (1985): Date Palms and Dates Pests. 1st Ed., University of Bassra, Ministry of Higher Education and Scientific Research, Baghdad, Iraq, 576 pp. (In Arabic).
- Hussain, S. and Aslam, N. A. (1970): Some observations on a beneficial Reduviid bug: *Amphibolus venator*, Klug (Fam. Reduviidae: Hemiptera). Agriculture Pakistan, 21 (1): 37-42.
- Hussein, A. E. (1990): Date varieties and palm height in relation to infestation with date stone beetle, *Coccotrypes dactyliperda* F. (Coleoptera: Scolytidae). Journal Annals of Agricultural Science, Moshtohor 1990 Vol. 28, No. (4): 2613-2622.
- Idder, M.A. and Pintureau, B. (2008): Efficacité de la coccinelle *Stethorus punctillum* (Weise) comme prédateur de l'acarien *Oligonychus afrasiaticus* (McGregor) dans les palmeraies de la région d'Ouargla en Algérie. Fruits, (63): 85-92.
- Innes, J. G. (1990): Ship Rat. The Handbook of New Zealand Mammals. King, C. M. (ed.) Oxford University Press: 206-225.
- Innes, J.; Hay, R.; Flux, I.; Bradfield, P.; Speed, H. and Jansen, P. (1999): Successful recovery of North Island kokako *Callaeas cinerea wilsoni* populations, by adaptive management. Biological conservation (87): 201-214.
- Jadhav, R. G.; Madane, N. P. and Kathamale, D. K. (1996): Record of soybean as a new host in India for citrus mealybug. Insect Environment, 2 (3): 90.
- Jalaluddin, S. M. and Sadakathulla, S. (1998): Effect of Atso oil emulsion sprays in the control of mealy bug *Maconellicoccus hirsutus* infesting guava fruits. Entomon 23: (2): 151-152.

References

- James, H. C. (1937): Sex ratios and the status of the male Pseudococcinae. Bulletin of Entomological Research, (28): 429-461.
- Jarjes, S. J.; Al-Mallah, N. M. and Abdulla, S. I. (1989): Insects and mites pests survey on rose-bay shrubs in Mosul region with some ecological and biological aspects of (*Nipaecoccus viridis* New.) and (*Parlatoria crypta* M) on rose-bay shrubs. Mesopotamia Journal of Agriculture, 21 (3): 29.
- Jaya, S.; Suresh, T.; Sobhitha Rani R. S. and Sreekumar, S. (2000): Evidence of seven larval instars in *Rhynchophorus ferrugineus* reared on sugarcane. Journal of Entomological Research (24): 27-31.
- Jepson, L. R.; Keifer, H. and Baker, E. W. (1975): Mites injurious to economic plants., University of California Press, Berkeley, 614 pp.
- Johnson, W. T. and Lyon, H. H. (1988): Insects that feed on trees and shrubs. Cornell University Press, 124 Roberts Place, Ithaca, New York 14850, U.S.A., 556 pp.
- Kaelin, P.; Zaugg, L.; Albertini, A. M. and Gadani, F. (1999): Activity of *Bacillus thuringiensis* isolates on *Lasioderma serricorne* (F.) (Coleoptera: Anobiidae). Journal of Stored Products Research, 35(2):145-158.
- Kairo M. T. K.; Pollard G. V.; Peterkin D. D. and Lopez, V. F. (2000): Biological control of the hibiscus mealybug, *Maconellicoccus hirsutus* Green (Hemiptera: Pseudococcidae) in the Caribbean. Integrated Pest Management Reviews, 5 (4): 241-254.
- Kamal, M. (1951): Biological control project in Egypt, with a list of introduced parasites and predators. Bulletin de la Societe Fouad I d'Entomologie, (35): 205-220.
- Kamali, K. and Taheri M. S. (1985): Morphological characteristics of *Ephestia elutella* (Hübner) (Lepidoptera; Pyralidae) and preliminary observations on its winter generation in Iran. Entomologie et Phytopathologie Appliquées, 52 (2): 89-97.
- Kane, E. and Ochoa, R. (2006): Detection and identification of the red palm mite *Raoiella indica* Hirst (Acari:Tenuipalpidae). USDA, ARS, Beltsville, MD, 6 pp.
- Kaya, H. K. and Gaugler, R. (1993): Entomopathogenic nematodes. Ann. Rev. Entomol. 38, 181-206. Gaugler, R., and Kaya, H. K. (Eds.). 1990. "Entomopathogenic Nematodes in Biological Control." CRC Press, Boca Raton, FL.
- Kerguelen, V. and Hoddle, M. S. (1999): Biological control of *Oligonychus perseae* (Acari: Tetranychidae) on avocado: II. Evaluation the efficacy of *Galendromus helveolus* and *Neosieulus californicus* (Acari: Phytoseiidae). International Journal of Acarology (25): 221-229.
- Khalaf, J. and Sokhansanj, M. (1993): Bioecological studies on orientalis yellow scale (*Aonidiella orientalis* New.) and its control by integrated methods in Fars province. Applied Entomology and Phytopathology, 60(1&2): ar 53-59, 11-12.
- Khan, M. A. (1983): Effectiveness of insecticides and repellents on stored-product pests. Anzeiger fur Schadlingskunde Pflanzenschutz Umweltschutz, 56 (2): 25-29.
- Khattak, S. U. K. and Jilani, G. (1984): Gamma radiation sensitivity of eggs of various ages of the red flour beetle, *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae. Pakistan Journal of Zoology (16): 21-24.

References

- Khawaja, A. H. and Akmal, M. (1971): Insect pests of date palm and their control. Date Palm Number, The Punjab Fruit Journal: 210-213.
- Khoualdia, O.; Rhouma A. and Hmidi, M. S. (1993): Contribution to the bio-ecological study of the white scale *Parlatoria blanchardi* Targ (Homoptera, Diaspididae) of date palm in Djerid (southern Tunisia). Annales de l'Institut National de la Recherche Agronomique de Tunisie, 66(1/2):89-108.
- Khoualdia, O.; Rhouma A.; Brun, J. and Marro, J. P. (1997): Biological control of white scale. Introduction of an exotic predator in the palm grove of Segdoud. Phytoma, 49 (494): 41-42.
- Khoualdia, O.; Rhouma, A. and Hmidi, M.S. (1993): Contribution to the bio-ecological study of the white scale *Parlatoria blanchardi* Targ. (Homoptera, Diaspididae) of date palm in Djerid (Southern Tunisia). Annales de l'Institut National de la Recherche Agronomique de Tunisie (66): 89-108.
- Kinawy, M. M. (1986): Coconut Rhinoceros Beetle, *Oryctes rhinoceros* L. (Coleoptera: Scarabidae) and its control. Ministry of Agriculture and Fisheries, Sultanate of Oman. ISBN (35/86), 24 pp. (In Arabic).
- Kinawy, M. M. (1987): Control of Coconut Rhinoceros Beetle, *Oryctes rhinoceros* L. (Coleoptera: Scarabidae) in South Oman. Bull. Fac. of Agric., Univ. of Cairo, 38 (1): 243-249.
- Kinawy, M. M. (1991): Biological control of the coconut scale insect (*Aspidiotus destructor* Sign., Homoptera: Diaspididae) in the southern region of Oman (Dhofar). Tropical Pest Management, 37 (4): 387-389.
- Kinawy, M. M. (1998): Biological control of onion thrips (*Thrips tapaci* Lind.) with the predatory mite (*Amblyseius cucumeris* (Oudemans), Acari: Phytoseiidae) on cucumber crops in the Sultanate of Oman. International Conference on Integrated Pest Management, Conference Proceedings, Sultan Qaboos University, 23-25 Feb., 1998, 67-73.
- Kinawy, M. M. (2004): Biological control of the coconut palm rhinoceros beetle, *Oryctes rhinoceros* L. Coleoptera: Scarabaeidae, by using *Rhabdionvirus oryctes* Hüger, in Sultanate of Oman. 1st Arab Conference of Applied Biological Pest Control, Cairo, Egypt, 5-7 April, 2004.
- Kinawy, M. M. (2007): Report of biological control of pomegranate fruit borer in Al-Jabal Al-Akhdar during 2007, Department of Plant Protection, Royal Gardens and Farms, Royal Court Affairs, Sultanate of Oman.
- Kinawy, M. M. and Hussein, M. H. (1987): Biological control of the citrus black fly (*Aleurocanthus woglumi* Ashby, Homoptera: Aleyrodidae) in South Oman. Bull. Fac. of Agric. Univ. of Cairo, 38 (1): 233-242.
- Kinawy, M. M.; Al-Waili, H. M. and Almandhari, A. M. (2008): Review of the successful classical biological control programs in Sultanate of Oman. Egyptian Journal of Biological Pest Control, 18 (1), 2008, 1-10.
- Klein, M. and Venezian, A. (1985): The dubas date tropiduchid, *Ommatissus binotatus lybicus*, a threat to date palms in Israel. Phytoparasitica (13): 95-101.
- Klotz, L.J. and Fawcett, H.S. (1932): Black scorch of the date palm caused by *Thielaviopsis paradoxa*. J. Agric. Res. (44): 155-166.
- Koppert, P. C. (1980): Bio logical control in glasshouses in the Netherlands. In: Proc. Int. Symp. IOBC/WPRS on Integrated control in Agriculture and Forestry, Vienna 8th – 12th October, 1979, 484 pp.

References

- Korunic, Z. and Hamel-Koren, D. (1985): The effect of deltamethrin and pirimiphos-methyl on pests of stored products. *Zastita Bilja*, 36 (4): 417-423.
- Kramer, V. L.; Garcia, R. and Colwell, A. E. (1988): An evaluation of *Gambusia affinis* and *Bacillus thuringiensis* var. *israelensis* as mosquito control agent in California wild rice fields. *J. Am. Mosquito Control Asso.* (4): 470-478.
- Kurian, C. (1963): Destructive pests of coconut other than *Oryctes rhinoceros*. Document, FAO, 1963. Technical working Party on Coconut, 1961, Bangkok.
- Lariviere, M. C. (1999): Cixiidae (Insecta: Auchenorrhyncha). Fauna of New Zealand, 40, ISBN 0-478-09334-9.
- Kurian, C.; Antony, J.; Abraham, V. A. and Natarajan. P. (1976): Pest management in coconut gardens. An integrated approach. *Indian Farming*, 7 (9): 31-35.
- LaHue, D. W. (1966): Evaluation of malathion, synergised pyrethrum and a diatomaceous earth as protectants against insects in sorghum grains in small bins. *Mark. res. rep.*, 781.
- Lai, P. Y. and Funasaki, G. Y. (1986): Introduction for biological control in Hawaii: 1983 & 1984. *Proceedings of the Hawaiian Entomological Society*, 26: 89-91.
- Langewald, J. and Schmutterer, H. (1995): Colour change in the desert locust, *Schistocerca gregaria* (Forsk.) (Orth., Cyrtacanthacridinae), induced by topical treatment with azadirachtin-enriched neem oil in semi-field trials in Benin at different population densities. *Journal of Applied Entomology*, 119 (3): 221-226.
- Langewald, J.; Kooyman, C.; Douro-Kpindou, O.; Lomer, C. J.; Dahmoud, A. O. and Mohamed, H. O. (1997): Field treatment of desert locust (*Schistocerca gregaria* Forsk!) hoppers in Mauritania using an oil formulation of the entomopathogenic fungus *Metarhizium flavoviride*. *Biocontrol Science and Technology*, 7 (4): 603-611.
- Langewald, J.; Ouambama, Z.; Mamadou, A.; Peveling, R.; Stolz, I.; Bateman, R.; Attignon, S.; Blanford, S.; Arthurs, S. and Lomer, C. (1999): Comparison of an organophosphate insecticide with a mycoinsecticide for the control of *Oedaleus senegalensis* (Orthoptera: Acrididae) and other Sahelian grasshoppers at an operational scale. *Biocontrol Science and Technology*, 9 (2): 199-214.
- Larivière, M. C. (1999): Cixiidae (Insecta: Hemiptera: Auchenorrhyncha). Fauna of New Zealand. Manaaki Whenua Press, 1999. ISBN 0-478-09334-9, Kote Aitanga Pepekeo Aotearoa, (40): 1-93.
- Larsen, T. B. (1980): Butterflies of Oman. John Bartholomew and Son Limited, UK, 80 pp. Lepesme, P. (1947): Les insectes des palmiers. Paris, Lechevalier.
- Larsen, T. B. (1985): Biogeographical aspects of Middle Eastern and Arabian butterflies. *Proceedings of the Symposium on the Fauna and Zoogeography of the Middle East*, 179-199.
- Larsen, T. B. and Nakamura, I. (1983): The Butterflies of East Jordan. *Entomologist Gazette*, 34: 135-208.
- Laurent, J. C. (1998): Acid rot. *Progrès Agricole et Viticole*, 115(1):7-9.
- Laville, E. (1973): Les maladies du dattier, pp. 95-108. In P. Munier, Ed.: *Le palmier dattier*. G.P. Maisonneuve & Larose, Paris, 221 pp.
- Le Pelley, R. H. (1968): *Pests of Coffee*. London and Harlow, UK: Longmans, Green and Co Ltd.

References

- Leefmans, S. (1920): De Palmsnuitkever (*Rhynchophorus ferrugineus* Olivier). Mededelingen van het Instituut voor Plantenziekten, No. 43, Department van Landbouw, Nigverheld en handel Batavia.
- Leeper, J. R. (1976): A review of the Hawaiian Coccinellidae. Proceedings of the Hawaiian Entomological Society, (22): 279-306.
- Lefkovitch, L. P. and Currie, J. E. (1967): Factors affecting adult survival and fecundity in *Lasioderma serricorne* (F.) (Coleoptera, Anobiidae). Journal of Stored Products Research, (3):199-212.
- Lepesme, P. (1947): Les insectes des palmiers. París: Paul Lechevalier, 904 pp.
- Lepigre, A. L. (1963): Essais de lutte sur l'arbre contre la pyrale des dattes (*Myelois ceratoniae* Zell., Pyralidae), Ann. Epiphytier, 14 (2): 85-101.
- Lever, R. J. W. (1969) : Pests of the Coconut Palm. FAO, Food and Agriculture Organization of the United Nations, Rome. Plant Production and Protection Series, No. (18), 190 pp.
- Li, Li-Ying (1984): Research and utilization of *Trichogramma* in China. In: Proceedings of the Chinese Academy of Sciences / United States National Academy of Sciences Joint Symposium on Biological Control of Insects, September 25-28, 1982, Beijing. Science Press, Beijing, 204-223.
- Li, Li-Ying. (1994): Worldwide use of *Trichogramma* for biological control on different crops: A survey. In biological control with egg parasitoids, eds. E. Wajnberg and S. A. Hassan, 37-53. Oxon, U.K., CAB International.
- Li, Yuezhong; Zeng-Rong Zhu; Ruiting Ju and Lian-Sheng Wang (2009): The Red Palm Weevil, *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae), Newly Reported from Zhejiang, China and Update of Geographical Distribution. Florida Entomologist (92): 386-397.
- Lindgren, D. L. and Vincent, L. E. (1953): Nitidulid beetles infesting California dates. Hilgardia (22): 97-118.
- Linsley, E. G. (1944): The date stone beetle in California and lower California. J. Econ. Ent. 36 (5): 804-805.
- Liotta, G.; Mineo, G. and Ragusa, S. (1976): On the current state of knowledge concerning certain arthropods injurious to citrus in Sicily. Bollettino dell'Istituto di Entomologia Agraria e dell'Osservatorio di Fitopatologia di Palermo, (10): 29-67.
- Lomer, C. J.; Bateman, R. P.; Dent D, DeGroote H.; Kooyman, C.; Langewald, J.; Johnson, D.; Peveling, R. and Thomas, M. B. (1999): Development of strategies for the incorporation of microbial pesticides into the integrated management of migratory pests. Agricultural and Forest Entomology, (1): 71-88.
- Louvet, J., Bulit, J., Toutain ,G. and Rieuf, P. (1970): Le Bayoud, Fusariose vasculaire du Palmier dattier. Symptômes et nature de la maladie. Moyens de lutte. Al Awamia, (35): 161-181.
- MacGowan, J. B. (1980): The American dagger nematode *Xiphinema americanum* Cobb, 1913. Nematology Circular, Division of Plant Industry, Florida Department of Agriculture & Consumer Services, 1980 No. 62: 2 pp.
- Madkour, M. (1979): Life cycle and biological control of the white scale. Training course of date palm pests and its control, 1979. Regional project for palm and dates research center in the Near East and North Africa. Baghdad, Iraq. (In Arabic).

References

- MAF (1994): Agricultural Research Annual Report, 1994, Ministry of Agriculture and Fisheries, Directorate General of Agricultural Research, Sultanate of Oman, 335 pp.
- Malencon, G. (1950): Le “Bayoud” maladie fusarienne du palmier-dattier en Afrique du Nord. Fruits d’outremer, (5): 279-289.
- Mani M. and Krishnamoorthy, A. (1996): *Aonidiella orientalis* (Newstead) (Diaspididae, Homoptera) and its natural enemies found on sapota, ber, custard apple and banana. Entomon, 21 (3/4): 273-274.
- Mani, M. (1989): A review of the pink mealybug - *Maconellicoccus hirsutus* (Green). Insect Science and its Application, 10 (2): 157-167.
- Mani, M. (1994): Effectiveness of the exotic encyrtid parasitoid, *Leptomastix dactylopii* How. in the control of mealybug, *Planococcus citri* (Risso) in guava orchards. Journal of Entomological Research, 18 (4): 351-355.
- Mani, M. and Krishnamoorthy, A. (2001): Suppression of *Maconellicoccus hirsutus* (Green) on guava. Insect Environment, 6 (4): 152.
- Mani, M. and Krishnamoorthy, A. (1999): *Maconellicoccus hirsutus* on acid lime in India. Insect Environment, 5 (2): 73-74.
- Marshal, K. J. (1970): Introduction of a new virus disease of the coconut rhinoceros beetle in Western Samoa. Nature 225: 288-289.
- Martin, H. (1958): Pests and diseases of date palm in Libya. FAO Plant Prot. Bull. 6 (8): 120-123.
- Martinez, A. P. (1966): FALSE SMUT OF PALMS. Plant Pathology Circular No. 43 Florida Department of Agriculture February Division of Plant Industry.
- Matsuura, H. (1993): Weevils associating with palms. Kobe Plant Protection 901, 46-47.
- Matthews, G. A. (1992): Pesticide application methods. Harlow, UK: Longman Scientific and Technical.
- McCoy, R. E. (1976): Comparative epidemiology of the lethal yellowing, kaincope and cadang-cadang diseases of coconut palm. Plant dis. Repr. (60): 498-502.
- McMurtry, J. A. and Croft, B. A. (1997): Life-styles of phytoseiid mites and their roles in biological control. Annual Review of Entomology (42): 291-321.
- McMurtry, J. A.; Huffaker, C. B. and Van de Vrie, M. (1970): Ecology of tetranychid mites and their natural enemies: a review. I. Tetranychid enemies: their biological characters and the impact of spray practices. Hilgardia 40, 331-390.
- Mekki, M. A. and Osman, M. H. (1991): Date Palm in Sultanate of Oman. Diwan of Royal Court, Muscat, Sultanate of Oman, 280 pp. (In Arabic).
- Menon, K. P. V. and Pandalai, K. M. (1960): Pests in the Coconut palm. A Monograph. Inrankulam, South India: Indian Central Committee, 261-265.
- Meyerdirk, D. E. and De Chi, L. W. (2003): “Models for Minimizing Risks of Dangerous Pests: The Pink Hibiscus Mealybug and Papaya Mealybug” Proceedings of the Caribbean Food Crops Society, Grenada, (39): 47-55.
- Meyerdirk, D. E.; Khasimuddin, S. and Bashir, M. (1988): Importation, colonization and establishment of *Anagyrus indicus* (Hym : Encyrtidae) on *Nipaecoccus viridis* (Hom.: Pseudococcidae) in Jordan. Entomophaga, 33 (2): 229-237.

References

- Michael, I. F. (1969): Studies on certain pests infesting date palm in U.A.R., Ph. D. Thesis, Fac. Of Agric., Ain Shams Univ., Cairo, Egypt.
- Michael, I. F. (1970): Economic importance and control of *Batrachedra amydraula* Meyr., the lesser date moth. Date growers's Inst. Rep. (47): 9-10.
- Michael, I. F. and Habib, A. A. (1971): Biology of *Batrachedra amydraula* Meyr., the lesser date moth. Date growers's Inst. Rep. 47: 5-8.
- Milner, R. J. (1970): The morphology and pathogenicity of Nosema whitei, a microsporidia pathogen of *Tribolium castaneum* (Hbst). Ph. D. Thesis, Newcastle upon Tyne Univ., U. K., 118 pp.
- Misra, H. P. and Bhatia, P. (1997): Biology of *Tribolium castaneum* strains resistant and susceptible to fenvalerate. Indian Journal of Entomology, 59 (2): 224-227.
- Misra, H. P. and Bhatia, P. (1998): Gamma radiation susceptibility of strains of *Tribolium castaneum* (Herbst) resistant and susceptible to fenvalerate. International Journal of Pest Management, 44(3):145-147.
- Mohaimed, A. G. (1978): Life cycle of the saw-toothed grain beetle, *Oryzaephilus surinamensis* (Coleoptera, Cucujidae) on dates in the middle and south of Baghdad. M. Sc. Thesis, Faculty of Agriculture, Baghdad University, Iraq.
- Mohamed, L. L. and EL-Haidary, H. (1965): A supplementary list of the phytophagous mites of Iraq. Ministry of Agric. Baghdad, Bull. 131.
- Mohammad, A. A.; Monir, M. M. and Hussein, A. E. (2004): Release of *Trichogramma evanescens* (West.) in Date Palm fields as an Eco-biological agent for suppression insect's infestation in Date Palms at El-Bahria Oases, Giza, Egypt. 1st Arab Conference of Applied Biological Pest Control, Cairo, Egypt, 5-7 April, 2004.
- Mokhtar, A. M. (2007): Integrated Pest Management of pomegranate Butterfly. Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 28 pp.
- Mokhtar, A. M.; Al-Nabahani, S. S. and Al-Tamimi, S. S. (2000a): Tracing the bio-ecology feature of *Ommatissus lybicus* De Berg. population on the light of honeydew production in situ. Research Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 31-42.
- Mokhtar, A. M.; Al-Nabahani, S. S. and Al-Tamimi, S. S. (2000b): Seasonal abundance of the fruit-stalk borer *Oryctes agamemnon* (Burn.) (Coleoptera: Scarabaeidae) in date palm plantations in Wadi Quriyat, Al-Dakhlyia Region. Research Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 44-46.
- Mokhtar, A. M.; Al-Nabahani, S. S. and Al-Tamimi, S. S. (2001a): Oviposition pattern of Dubas bug *Ommatissus lybicus* De Berg. On date palm. Research Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 25-40.
- Mokhtar, A. M.; Al-Nabahani, S. S. and Al-Tamimi, S. S. (2001b): Estimation of honeydew production of different stages of *Ommatissus lybicus* De Berg. (Homoptera: Tropiduchidae) in vitro. Research Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 14-24.

References

- Mokhtar, A. M.; Al-Nabahani, S. S. and Al-Tamimi, S. S. (2001c): Using the production of honeydew as indication to estimate the population of Dubas Bug *Ommatissus lybicus* De Berg. In the date palm grooves in Oman. Research Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 3-13.
- Mokhtar, A. M.; Al-Nabahani, S. S. and Al-Tamimi, S. S. (2001d): Collection and identification of new insect at ARS, Jimah, *Perindus binudatus* Emeljanov (Hemiptera: Cixiidae). Research Annual Report of Entomology, Jimah Research Station, Ministry of Agriculture and Fishers, Sultanate of Oman, 88-92.
- Mokhtar, A. M.; Lwatyia, M. H. and Al-Abd, M. (1991): Light Trap Monitoring of Insects. Agricultural Research Annual Report, Directorate General of Agricultural Research, Rumais, Sultanate of Oman, 206-213.
- Monciero, A. (1947): Etude comparée sommaire des différents types de culture du palmier dattier en Algérie. Fruits (2): 374-382.
- Mondal, K. (1985): Response of *Tribolium castaneum* larvae to aggregation pheromone and quinones produced by adult conspecifics. International Pest Control, 27 (3): 64-66.
- Montaigne, Mde. and Maouloud, F. A. (1986): The sanitary protection of palm trees in Mauritania. Phytoma, No. 38: 41-45.
- Monty, J. (1974): Teratological effects of the virus *Rhabdionvirus oryctes* on the coconut palm rhinoceros beetle *Oryctes rhinoceros* (L.). Bull. Entomol. Res. (64): 633-636.
- Moreno, D. S.; Fagerlund, J. and Ewaet, W. H. (1984): Citrus mealybug *Planococcus citri* (Homoptera: Pseudococcidae): behaviour of males to sex pheromone in laboratory and field. Annals of the Entomological Society of America, 77 (1): 32-38.
- Moutia, L. A. (1958): Phytophagous Acarina and their predators in Mouritius. Bull. Ent. Res., (49): 59-75.
- Munier, P. (1955): Le palmier dattier en Mauritanie. Ann. Inst. Fruits et Agrumes Coloniaux 12, 66 pp.
- Murphy, S. T. and Biscoe, B. R. (1999): The red palm weevil as an alien invasive: biology and the prospects for biological control as a component of IPM. Biocontrol News and Information, (20): 35-46.
- Murphy, S. T. and Briscoe, B. R. (1999): The red palm weevil as an alien invasive: biology and the prospects for biological control as a component of IPM. Biocontrol News and Information, Vol. 20 No. 1, 35N-46N.
- Muthu, M.; Rajendran, S.; Krishnamurthy, T. S.; Narasimhan, K. S.; Rangaswamy, J. R.; Jayaram, M. and Majumder, S. K. (1984): Ethyl formate as a safe general fumigant. Controlled atmosphere and fumigation in grain storages. Proceedings of an international symposium 'Practical aspects of controlled atmosphere and fumigation in grain storages' held from 11 to 22 April 1983 in Perth, Western Australia, 369-393.
- Myres, L. E. (1932): Two economic greenhouse mealybugs of Mississippi. Journal of Economic Entomology, 25:891-896.
- Nixon, R. W. (1932): Observations on the occurrence of blacknose. Ann. Date Growers' Instit. (9): 3-4.
- Nixon, R. W. (1951): Date culture in the United States. U.S.D.A. Circ. 728 Revised. 57 pp., illus.
- Nixon, R. W. (1954): Date culture in Saudi Arabia. Ann. Date grower's Instit. (31): 15-20.

References

- Nixon, R. W. and Carpenter, J. B. (1978): Growing Dates in the United States. US Department of Agriculture, Washington, DC., Bull. 107.
- Oatman, E. R. and Platner, G. R. (1985): Biological control of two avocado pests. Calif. Agric., 39 (11/12): 21-23.
- Orton-Williams, K. J. (1972): *Meloidogyne javanica*. CIH Descriptions of Plant Parasitic Nematodes, Set 1, No. 3. Wallingford, UK: CAB International.
- Osman, A. M. and Abdel Redha, A. H. (1989): Palm in Kuwait. Department of Agricultural Extension, Plant Wealth Management, Public Authority for Agriculture Affairs and Fish Resources.
- Pacheco, I. A.; Santori, M. R. and Taylor, R. W. D. (1990): Survey of phosphine resistance in stored grain insect pests in the state of Sao Paulo. Coletânea do Instituto de Tecnologia de Alimentos, 20 (2):144-154.
- Padi, B. (1996): Parthenogenesis in mealybugs occurring on cocoa, pp. 242-248. In Proceedings of the First International Cocoa Pests and Diseases Seminar, Accra, Ghana, Nov. 1995.
- Panis, A. (1979): Mealybugs (Homoptera, Coccoidea: Pseudococcidae) within the framework of integrated control in Mediterranean citrus-growing. Revue de Zoologie Agricole et de Pathologie Vegetale, 78 (3): 88-96.
- Panis, A. and Brun, J. (1971): Biological control tests against three species of *Pseudococcidae* (Homoptera, Coccoidea) in greenhouses. Revue de Zoologie Agricole et de Pathologie Vegetale, 70 (2): 42-47.
- Paulin-Mahady, A. E.; Harrington, T. C. and McNew, D. L. (2002): Phylogenetic and taxonomic evaluation of Chalara, Chalaropsis, and *Thielaviopsis anamorphs* associated with Ceratocystis. Mycologia (94): 62-72.
- Pereau-Leroy, P. (1958): Le plamier- dattier au Maroc. Min. Agric. Maroc, Serv. Rech. Agron. Et Inst. Francais Rech. Fruit outer mer Rabat, 142 pp.
- Peterson, A. (1960): Larvae of insects, part I and II. Ohio State Univ.Columbus, Ohio, U.S.A.
- Pollard, G.V., (1995): Pink or hibiscus mealybug in the Caribbean. CARAPHIN News, 12: 1-2.
- Prevett, P. F.; Benton F. P.; Hall, D. R.; Hodges, R. J. and Serodio, R. S. (1989): Suppression of mating in *Ephestia cautella* (Walker) (Lepidoptera: Phycitidae) using microencapsulated formulations of synthetic sex pheromone. Journal of Stored Products Research, 25 (3): 147-154.
- Price, L. A. and Mills, K. A. (1988): The toxicity of phosphine to the immature stages of resistant and susceptible strains of some common stored product beetles, and implications for their control. Journal of Stored Products Research, 24 (1): 51-59.
- Price, R. E.; Bateman, R. P.; Brown, H. D.; Butler, E. T. and Müller, E. J. (1997): Aerial spray trials against brown locust (*Locustana pardalina* Walker) nymphs in South Africa using oil-based formulations of Metarhizium flavoviride. Crop Protection, 16 (4): 345-351.
- Pritchard, A. E. and Baker, E. W. (1955): A revision of the spider mite family Tetranychidae. Pacific Coast Ent. Soc. Memoirs Series, Vol. 2.
- Prozell, S. and Scholler, M. (1997): The insect fauna of an industrial bakery applying biological control with *Trichogramma evanescens* Westwood instead of synthetic chemical insecticides. Proceedings

References

- of the German Society for General and Applied Entomology, Bayreuth, Germany, 18-22 March 1997. Mitteilungen der Deutschen Gesellschaft fnr Allgemeine und Angewandte Entomologie, 11 (1-6): 293-296.
- Rabb, R. L. (1972): Principles and concepts of pest management. In implementing practical pest management strategies. Proceedings of a national extension pest-management workshop. Purdue University, Lafayette, Indiana, 6-29.
- Rajagopal, D. and Krishnamoorthy, A. (1996): Bionomics and management of oriental yellow scale, *Aonidiella orientalis* (Newstead) (Homoptera: Diaspididae): an over view. Agricultural Reviews (Karnal), 17 (3/4): 139-146.
- Ramakers, P. M. J. (1978): Possibilities of biological control of *Thrips tabaci* Lind. (Thysanoptera: Thripidae) in glasshouses. Med. Fac. Landbouww. Rijksuniv. Gent (43): 463-469.
- Rangaswamy, J. R. and Sasikala, V. B. (1991): Sex pheromone properties of Z-2,nonenyl propionate isolated from virgin females of red flour beetle *Tribolium castaneum* (Coleoptera). Indian Journal of Experimental Biology, 29 (3): 263-266.
- Rao, Y. R. (1921): A preliminary list of insect pests of Iraq. Department of Agriculture of Iraq. Memoir No. 7, Time Press, Bombay.
- Reddy, K. B., Prakasan, C. B., Bhat, P. K. and Kumar, A. C. (1992): Establishment of *Leptomastix dactylopii* How. (Hym.: Encyrtidae) in Karnataka for control of *Planococcus citri* (Risso) (Hom.: Pseudococcidae) of coffee. Journal of Coffee Research, 22 (1): 37-44.
- Reginald, C. (1973): Principal Insect Pests. In: Coconuts. Tropical Agriculture Series, London, UK: Longmans.
- Reisi, P. R.; Pedro Neto, M.; Franco, R. A. and Teodoro, A. V. (2003): Control of *Brevipalpus phoenicis* (Geijskes, 1939) and *Oligonychus ilicis* (McGregor, 1917) (Acari: Tenuipalpidae, Tetranychidae) in coffee plants and the impact on beneficial mites: I-Abamectin and emamectin. Ciênc. agrotec. [online]. 2004, Vol. 28, No. (2), 269-281.
- Rieuf, P. (1968): La maladie des tâches brunes du palmier dattier. Al Awamia, (26):1-24.
- Rizk, G. N.; El-Badry, E. and Hafez, S. M. (1979): The effectiveness of predacious and parasitic mites in controlling *Tribolium confusum* Duv. Mesopotamia Journal of Agriculture, 14 (2): 167-182.
- Rodriguez, J. G. and Rodriguez, L. D. (1987): Nutritional Ecology of Stored Product and House Dust Mites. In: F. Slansky, J.G. Rodruquez (Eds). Nutritional Ecology of Insects, Mites, Spiders, and Related Invertebrates. Wiley., 345-368.
- Rotundo, G. and Tremblay, E. (1974): Studies on a sexual pheromone or *Planococcus citri* (Risso) (Homoptera, cocoidea). II. Extraction. Bollettino del Laboratorio di Entomologia Agraria 'Filippo Silvestri', Portici, 31: 121-131.
- Roy, M.; Brodeur, J. and Cloutier, C. (2003): Effect of temperature on intrinsic rates of natural increase (r_m) of a Coccinellid and its spider mite prey. BioControl (48): 57-72.
- Saaidi, M. (1979): Contribution à la lutte contre le Bayoud, *Fusariose vasculaire* du palmier. Thèse d'Université. Université de Dijon, 140 pp.
- Saaidi, M. (1992): FAO Report on date palm protection and training activities. Ministry of Agriculture & Fishers. Sultanate of Oman, 121 pp.

References

- Sachs, G. (1967): Sur la présence d'Omphalia sp. Bliss dans une palmeraie Mauritanienne. *Fruits* 22: 497-501.
- SAEED, A. (1972): Date palm number. *The Punjab Fruit Journal*. vol 33/34: No. 114-119.
- Sadof, C. (1995): Know Your Friends: Mealybug Destroyer, Midwest Biological Control News Online, II: 5.
- Sass, B. D. and Wyatt Hoback, W. (2006): Effects of Temperature and Humidity on Grain Mite, *Acarus siro*, Survival, ESA / NCB Meeting.
- Sathiamma, B.; Abraham, V. A. and Kurian, C. (1982): Integrated pest management of the major pests of coconut. *Indian Coconut Journal*, 12 (6/9): 27-29.
- Sauer, D. B. (1992): Storage of Cereal Grains and Their Products. 4th edition. St Paul, USA: American Association of Cereal Chemists, Inc.
- Sayed, M. T. (1942): Contribution to the knowledge of acarina in Egypt: The genus *Raoiella* Hirst (Pseudotetranychinae:Tetranychidae). *Bull. Soc. Fouad Ier DEntomologie*. 26: 81-91.
- Schmutterer, H.; Baumgart, M.; Freisewinkel, D.; Langenwald, J. and Nicol, C. M. Y. (1993): The effects of neem oil and other neem products on nymphs and resulting adults of *Schistocerca gregaria*, *Nomadacris septemfasciata*, *Locusta migratoria migratorioides*, and *Zonocerus variegatus*. *Journal of Applied Entomology*, 116 (2): 178-186.
- Sen Selvan, P.; Grewal, S.; Gauglar, R. and Tomalak, M. (1994): Evaluation of steinernematid nematodes against *Popillio japonica* larvae: species, strains, and rinse after application. *Journal of Economic Entomology* 87 (3): 605-609.
- Shamseldean, M. M. and Abdel-Gawad, M. M. (1994): Laboratory evaluation of Heterorhabditid nema-
- todes for control of the palm weevil. *Egyptian Journal of Applied Sciences*, 9 (3): 670-679.
- Sharaf, N. S. (1996): Importance of life tables for determining proper timing and frequency of insecticide application in controlling the spherical mealybug, *Nipaecoccus viridis* (Newstead) (Homoptera: Pseudococcidae). *Dirasat. Agricultural Sciences*, 23 (2): 103-110.
- Sher, S. A. (1966): Revision of the Hoplolaiminae (Nematoda) VI. *Helicotylenchus Steiner, 1945. Nematologica*, (12):1-56.
- Sher, S. A. and Allen, M. W. (1953): Revision of the genus *Pratylenchus* (Nematoda: Tylenchidae). *Univ. Calif. Publ. Zool.*, (57): 441-470.
- Shorrocks, B. (1972): *Drosophila*. Ginn & Company Limited: London. *Drosophila*, 144 pp.
- Shtakel'berg, A. A. (1970): Family Drosophilidae. In: Bienko G Ya Bei, ed. Keys to the insects of the European part of the USSR. *Opredeliteli po Faune SSSR*, 103 (5): 390-399.
- Siddiqi, M. R. (1972): *Helicotylenchus dihystera*. C.I.H. Descriptions of Plant Parasitic Nematodes, Set 1: 9. Wallingford, UK: CAB International, 3 pp.
- Siddiqi, M. R. (1973): *Xiphinema americanum*. C.I.H. Descriptions of Plant-parasitic Nematodes, Set 2 (No. 29): 4 pp.
- Simmonds, H. W. (1941): Biological control of the rhinoceros beetle, *Oryctes rhinoceros*. Department of Agriculture, Fiji. Bulletin No. 21.
- Simmonds, H. W. (1949): On the introduction of *Scolia ruficornis* F. into Western Samoa for the control of *Oryctes rhinoceros*. *Bull. Ent. Res.*, No. (40): 445-446.

References

- Singh, M. P. and Ghosh, S. N. (1970): Studies on *Maconellicoccus (Phenacoccus) hirsutus* Gr. causing 'bunchy top' in mesta. Indian Journal of Science and Industry, A, 4 (2): 99-105.
- Singh, S. P.; Rao, N. S.; Ramani, S. and Poorani, J. (2001): Research highlights 2000-2001. Bangalore, India, 21 pp.
- Smirnoff, W. A. (1957): La cochenille du palmier-dattier (*Parlatoria blanchardii* Targ.) en Afrique du Nord. Comportement, importance économique, prédateurs et lutte biologique. Entomophaga, (2): 1-99.
- Smith, D.; Papacek, D. F. and Murray, D. A. H. (1988): The use of *Leptomastix dactylopis* Howard (Hymenoptera: Encyrtidae) to control *Planococcus citri* (Risso) (Hemiptera: Pseudococcidae) in Queensland citrus orchards. Queensland Journal of Agricultural and Animal Sciences, 45 (2): 157-164.
- Smith, D.; Smith, I. W. and Clements, P. (2003): Fusarium Wilt of Canary Island Date Palm. Forests Fact Sheet, Victoria, Australia: Department of Sustainability and Environment.
- Smith, E. H. (1972): In implementing practical pest management strategies. Proceedings of a national extension pest-management workshop. Purdue University, Lafayette, Indiana.
- Smith, K. G. V. (1989): An introduction to the immature stages of British flies: Diptera larvae, with notes on eggs, puparia and pupae. Handbooks for the Identification of British Insects London, UK: Royal Entomological Society.
- Smith, M. T.; Georgis, R.; Nyczepir, P. and Miller, R. W. (1993): Biological control of the pecan weevil, *Curculio caryae* with entomopathogenic nematodes. Journal of Nematology, 25 (1): 78-82.
- Smith, S. M. (1996): Biological control with *Trichogramma*: Advances, success and potential of their use. Ann. Rev. Entomol, (41): 375-406.
- Smith-Meyer, M. K. P. (1987): African Tetranychidae (Acari: Prostigmata) - with reference to the world fauna. Entomology Memoir, Department of Agriculture and Water Supply, Republic of South Africa, (No. 69): 77-78, 80-82.
- Snowdon, A. L. (1990): A colour Atlas of Post-Harvest Diseases & Disorders of Fruits & Vegetables. Volume 1: General Introduction and Fruits. University of Cambridge. Publishing Ltd, 2-16 Torrington Place, London WC1E 7LT, England, 302 pp.
- Stansly, P. A. (1984): Introduction and evaluation of *Chilocorus bipustulatus* (Coleoptera: Coccinellidae) for control of *Parlatoria blanchardi* (Homoptera: Diaspididae) in date groves of Niger. Entomophaga, 29 (1): 29-39.
- Stickney, F. S.; Barnes, D. W. and Simmons, P. (1950): Date Palm Insects in the United States. Circular No. 846, US Department of Agriculture, Washington, DC.
- Stojanovic, T. and Kosovac, V. (1974): Degree of insect infestation of stored sunflower in Banat in the period 1966-1974. Zastita Bilja, 25 (128/129): 183-192.
- Strand, L. L. (1993): Integrated Pest Management for Strawberries. Pub. 3351. University of California. 15 pp.
- Suzuki, T. (1985): Presence of another aggregation substance(s) in the frass of the red flour beetles, *Tribolium castaneum* (Coleoptera: Tenebrionidae). Applied Entomology and Zoology, 20 (1): 90-91.
- Swaminathan, R. and Verma, S. K. (1991): Studies on the incidence of date palm scale, *Parlatoria*

References

- blanchardi* (Targ.) in western Rajasthan. Entomon, 16 (3): 217-221.
- Takagi, S. (1970): Diaspididae of Taiwan based on material collected in connection with the Japan. U. S. Co-operation Science programme, 1965 (Homoptera : Coccoidea). Insecta Matsumurana, 33 (2): 53-67.
- Talhouk, A. S. (1983): The present status of date palm pests in Saudi Arabia. Proceedings of the First Symposium on the Date Palm in Saudi Arabia, 23-25 March 1982, King Faisal Univ., Al-Hassa, Kingdom of Saudi Arabia, 432-437.
- Talhouk, A. S. (1991): On the management of the date palm and its arthropod enemies in the Arabian Peninsula. Journal of Applied Entomology 111, 514-520.
- Tanigoshi, L. K. and McMurtry, J. A. (1977): The dynamics of predation of *Stethorus picipes* (Coleoptera: Coccinellidae) and *Typhlodromus floridanus* on the prey *Oligonychus punicae* (Acarina: Phytoseiidae, Tetranychidae). Hilgardia (45): 237-288.
- Tauber, C.A. (1991): Order Neuroptera. In Immature Insects. Vol. 2, F. W. Stehr, ed. Dubuque, IA: Kendall Hunt. 126-143.
- Tawfik, M. F. S. (1997): Biological control of agricultural pests. Academic Bookshop. Cairo, Egypt pp. 757.
- Tawfik, M. F. S.; Awadallah, K.T. and Abdella, M. M. H. (1982): Effect of prey on various stages of the predator *Allaeocranum biannulipes* (Montr. et Sign.) (Hemiptera; Reduviidae). Bulletin de la Société Entomologique d'Egypte, No. 64:251-258.
- Tawfik, M. F. S.; Awadallah K. T.; El-Husseini, M. M. and Ibrahim, A. M. A. (1986a): Effect of temperature on the life-cycle of the predator *Termatophyllum insigne* Reuter (Hemiptera: Termatophylidae). Bulletin de la Société Entomologique d'Egypte, No. (66): 43-49.
- Tawfik, M. F. S., Awadallah K. T.; El-Husseini, M. M. and Ibrahim, A. M. A. (1986b): Effect of temperature and relative humidity on the biocycle of *Xylocoris sordidus* (Reuter) (Anthocoridae, Hemiptera). Archiv für Phytopathologie und Pflanzenschutz, 22 (2): 119-129.
- Taylor, R. W. D. (1991): Resistance to grain fumigants and future prospects for their use. Pesticide Outlook, 2 (2): 22-24.
- Temerak, S. A. and Sayed, A. A. (2001): Ovi-larvicidal activity of spinosad in comparison to *Bacillus thuringiensis* subsp kurstaki for the control of *Virachola livia* (Klug.) on date palm trees in the field, New Valley, Egypt. Assiut J. of Agric., Sci., Vol. 32, No. 4, 2001.
- The Natural History Museum. (2002): Ref. No. ENQ 2002/734, 11 September 2002. Identification of *Oligosita* sp. from Oman.
- Thomas, D. L. (1974): Possible link between declining palm species and lethal yellowing of coconut palms. Proc. Fla. State Hort. Soc. 87: 502-504.
- Tourneur, J. C. and Lecoustre, R. (1975): Cycle De Developpement Et Tables De Vie de *Parlatoria blanchardi* Targ. (Homoptera: Diaspididae) Et de son predateur exotique En Mauritanie, *Chilocorus bipustulatus* l. var. *Iranensis* (Coleóptera: Coccinellidae). Fruits (30): 481-497.
- Toutain, G. (1965): Note sur l'épidémiologie du Bayoud en Afrique du Nord – Al Awamia, 15: 37-45.
- Toutain, G. (1970): Observations sur la progression d'un foyer actif de Bayoud dans une plantation régulière de palmier dattier. Al Awamia, (35): 155-160.

References

- Toutain, G. (1973): Lutte contre le Bayoud. 1. Reconstitution de la palmeraie bayoudee au Macroc. Al Awamia, (48): 115-146.
- Toutain, G. (1976): The date palm: Culture and production. Al Awamia, Rabat No. (25): 23-151.
- Toutain, G. and Louvet, J. (1972): Résistance au Bayoud dans les variétés de palmier dattier, In H. Fargi and J. M. Sonpault eds., le palmier dattier et sa Fusarios vasculaire (Bayoud) DRA (Maroc) INRA (France).
- Urban, A. and Schmidt, H. (1993): Moth pests on sultanas controlled biologically. Plant Protection News, No. (31): 8-9.
- Uvarov, B. P. (1921): A revision of the genus *Locusta*, L. (=Pachytalus, Fieb.), with a new theory as to the periodicity and migrations of locusts. Bull. Ent. Res., 12: 135-163.
- Uvarov, B. P. (1928): Locusts and grasshoppers. London, Imperial Bureau of Entomology, 352 pp.
- Uvarov, B. P. (1966): Grasshoppers and locusts. A handbook of General Acridology, I, II. Cambridge Univ. Press, Cambridge, 481 pp.
- Vanderplank, F. L. (1958): The assassin bug, *Platymeris rhadamanthus* Gerst., a usful predator of Oryctes boas and O. rhinoceros. J. ent. Soc. South Africa, II, 21 (2): 309-314.
- Vidyasagar, P. S. P. V.; Al-Saihati, A. A.; Al-Mohanna, O. E.; Subbei, A. I. and Abdul Mohsin, A. M. (2000): Management of red palm weevil *Rhynchophorus ferrugineus* Oliv., a serious pest of date palm in Al Qatif, Kingdom of Saudi Arabia. Journal of Plantation Crops, 28 (1): 35-43.
- Viggiani, G. (2008): Identification of Dubas bug egg parasitoid *Pseudoligosita babylonica* from Oman (Personal Contact).
- Wajnberg, E. and Hassan, S. A. (1994): Biological control with egg parasitoids. CAB International, International Institute of Biological Control, Silwood Park, Buckhurst Road, Ascot, Berkshire SL5 7TA, UK: 286 pp.
- Walker, D. H. and Pittaway, A. R. (1987): Insects of Eastern Arabia. Illustrated by A. J. Walker. Published by Macmillan Publishers Ltd, 175 pp.
- Waller, J. H. and Bridge, J. (1978): Plant diseases and nematodes in the Sultanate of Oman. PANS 24: 313-326.
- Wattanapongsiri, A. (1966): A Revision of the Genera *Rhynchophorus* and *Dynamis* (Coleoptera: Curculionidae). Bangkok, Thailand. Dept. Agric. Sci. Bull. 1-329.
- Welbourn, C. (2006): Red Palm Mite *Raoiella indica* (Acari: Tenuipalpidae). Pest Alert. DPI-FDACS; 4pp. (<http://www.doacs.state.fl.us/pi/enpp/ento/r.indica.html>).
- West, C. P.; Izekor, E.; Oosteruis, D. M. and Robbins, R. T. (1988): The effect of *Acremonium coenophialum* on growth and nematode infestation of tall Fescue. Plant and Soil, 112: 3-6.
- Willcocks, F. C. (1914): The date-stone beetle. Bull. Ent. d'Egypte, (6): 37-39.
- Williams, D. J. (1996): A brief account of the hibiscus mealybug *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), a pest of agriculture and horticulture, with descriptions of two related species

References

- from southern Asia. Bulletin of Entomological Research, 86 (5): 617-628.
- Williams, D. J. and Watson, G. W. (1988): The scale insects of the tropical South Pacific Region. Part 1. The armoured scales (Diaspididae). Wallingford, UK: CAB International.
- Wilps, H.; Nasseh, O.; Rembold, H. and Krall, S. (1993): The effect of *Melia volkensii* extracts on mortality and fitness of adult *Schistocerca gregaria* Forskål (Orth., Cyrtacanthacrinae). Investigations conducted under natural conditions in *S. gregaria* recession areas in the southern Tamesna Desert (Republic of Niger). Journal of Applied Entomology, 116 (1): 12-19.
- Woets, J. (1973): Integrated control in vegetable under glasshouse in the Netherlands. Bull. O. L. L. B. S. R. O. P. (4): 26-31.
- Yablokov-Khnzoryan, S. M. (1978): Beetles of the tribe Laemophloeini (Coleoptera, Cucujidae) in the Soviet fauna. Communication 2. Entomological Review, (57): 237-249.
- Young, E. C. (1974): The epizootiology of two pathogens of the coconut palm rhinoceros beetle. J. Invertebr. Pathol., (24): 82-92.
- Yousif, A. and Al-Bukiri, S. (2003): Effect of Irrigation on Within – Grove Distribution of Red Palm Weevil, *Rhynchophorus ferrugineus*. Sultan Qaboos University, Agricultural and Marine Sciences, 8 (1): 47-49.
- Zaher, M. A.; Wafa, A. K. and Yousef, A. A. (1969): Biological studies on *Raoiella indica* Hirst and *Phyllozetranychus aegyptiacus* Sayed infesting date palm trees in U.A.R. (Acarina: Tenuipalpidae). Zeitschrift für angewandte Entomologie.
- Zaher, M. A.; Wafa, A. K. and Yousef, A. A. (1971): Biology of *Brevipalpus phoenicis* (Geijskes), in Egypt (Acarina: Tenuipalpidae). Journal Bulletin de la Societe Entomologique d'Egypte, publ. 1971 Vol. (54): 177-183.
- Zaid, A. (1999): African Palm Weevil *Rhynchophorus phoenicis* F. Attack on Date Palm in the Republic of South Africa and Zimbabwe. Middle East Red Palm Weevil Workshop, 26-29 January, 1999, Cairo-Egypt.
- Zaki, F. N. (1989): Rearing of two predators, *Orius albidipennis* (Reut.) and *Orius laevigatus* (Fieber) (Hem., Anthocaridae) on some insect larvae. Journal of Applied Entomology, 107 (1): 107-109.
- Zelazny, B. (1977): *Oryctes rhinoceros* population and behaviour influenced by a baculovirus. Journal of Invertebrate Pathology 29: 210-215.
- Zerova, M. D. and Fursov, V. N. (1989): A catalogue of species of the genus *Trichogramma* Westwood of the world (Hymenoptera: Trichogrammatidae). Institute of Zoology of the Science Academy of the Ukrainian SSR. Publ. 89 (4), 52 pp.

Index

A

Abelmoschus esculentus 54
Acacia catechu 179
Acacia farnesiana 179
Acacia horrida 179
Acacia lasiopetala 179
Acacia mellifera 179
Acacia moderata 179
Acacia nilotica 179
Acacia raddiana 132
Acacia spalicigora 179
Acacia spp. 177
Acacia tortilis 54
Acari 170, 34, 205, 210, 211, 213, 214, 215, 219, 202, 217
Acarina 41, 185
Acaropsellina docta Berlese 159
Acoelorrhaphes wrightii 238, 255
Acomys dimidiatus Thomas 286
Acrididae 131, 136
Adeleidae 159
Adelina tribolii 159
Adiheteothrips jambudvipae Ramok 142
Adoxophyes orana 351
Adoxophyes orana Fischer von Roesl 351
Aerocerus fasciculatus 195
Agaricales 245
Agelenidae 340
Agrobacterium tumefasciens 363
Aiphanes lindeniana 255
Albizzia julibrissin 54
Alcterogystia cadamiae (Moore) 326
Allagoptera arenaria 255
Alternaria 352
Alternaria alternata (Fr) Keissler 247
Alternaria sp. 187
Amblyseius caudatus Berlese 213
Amblyseius channabasavanni Gupta & Daniel 213
Amblyseius cucumeris (Oudemans) 340
Amblyseius largoensis (Muma) 213, 217
Amblyseius longispinosus Evans 213
Amblyseius mackenziei Schuster & Pritchard 340
Amblyseius spp. 339
Ameroseiidae 104

Ameroseius sp. 104
Amitermes desertorum (Deseneux) 140
Amitermes gallagheri Chhotani & Bose 140
Amitermes stephensi Harris 140
Amitermes vilis Hagen 140
Amorbia cuneana (Walsingham) 345
Amphibolus venator (Ashmead) , 159
Amphisphaeriaceae 237
Anacridium melanorhodon arabafrum 136
Anagrapha falcifera 351
Anagyrus agraensis Saraswat 327, 64
Anagyrus bohemanni Westwood 64
Anagyrus dactylopis (Howard) 327, 348
Anagyrus diversicornis Howard 64
Anagyrus indicus Shafee 64
Anagyrus kamali 58
Anagyrus kamali Moursi 58
Anagyrus mirzai Agarwal 327, 348
Anagyrus pseudococcii (Girault) 61
Ananas comosus 116
Angulitermes arabiae Chhotani & Bose 140
Anobiidae 162
Anthocoridae 147, 220
Anticarsia gemmatalis 351
Anystidae 338
Aonidiella orientalis 331
Aonidiella orientalis (Newstead) 44
Apanteles sp. 79
Aphelinida 278, 279
Aphelinidae 41, 46
Aphis gossypii (Glover) 333
Aphis punicae (Passerini) 333
Aphytis lingnanensis Compere 327, 46
Aphytis melinus DeBach 46
Aphytis mytilaspidis Lebaron 41
Aphytis phoeniceis Debaels & Rosen 41
Aprostocetus sp. 327
Arachnida 340
Araneae 327, 340
Araneus diadematus Clerch 341
Araneus spp. 341
Archenomus arabicus Ferriere 349
Archirileya femorata (Boucek) 69
Areca catechu 44, 82

Arecastrum romanoffianum (Chamisso) Beccari 232
Arega engleri 255
Arega pinnata 255
Arena pinnata 82, 238
Arenipes sabella Hampson 77
Armascirus taurus (Kraemer) 213
Arthropoda 338
Ascidae 170
Aspergillus niger Van Tieghem 248
Aspergillus sp. 187
Aspidiotus destructor Sign. 331
Asterolecaniidae 50
Asterolecanium phoenicis (Ram. - Rao) 50, 320
Asterolecanium pustulans (Cockerell) 50
Auchenorrhyncha 253
Autographa californica 351
Azadirachta indica 44, 132, 135

B

Bacillus thuringiensis 163
Bacillus thuringiensis subsp. *kurstaki* (Btk) 351
B. t. subsp. *aizawai* 351
B. t. subsp. *japonensis* strain *buibui* 351
B. t. subsp. *tenebrionis* 351
Bactris gasipaes 255
Baculovirus oryctes 113
Bandicota bengalensis (Gray) 286
Bandicota (*Nesokia*) *indica* 286
Batrachedra amydraula Meyrick 72
Beauveria bassiana (Bals.-Criv.) Vuill. 99, 104, 135, 352
Beauveria brongniartii 352
Bemisia tabaci 352
Bethylidae 76
Bismarckia nobilis 238
Blattisocius tarsalis Berlese 170
Blepharopsis mendica Fabricius 337
Blighia sapida 50
Bocchus hyalinus Olmi 34
Bochartia sp. 34
Bombycillidae 307
Borassus flabellifer 255
Bostrichidae 120
Botrydiplodia theobromae (Pat.) 230
Botryosphaeriaceae 235
Botryosphaeraiales 235
Brachymeria aegyptiaca (Masi) 181

Bracon brevicornis Wesmael 76, 178, 202
Bracon (*Habrobracon*) *hebetor* Say 76, 202
Braconidae 170, 178, 202
Brevipalpus phoenicis (Geijskes) 215
Buprestidae 122, 195
Burkholderia cepacia 352
Butia capitata 238

C

Cadra cautella (Walker) 202
Cadra cautella Walker 196
Cadra (*Ephestia*) *calidella* Guenée 171
Cadra (*Ephestia*) *cautella* (Walker) 167
Cadra (*Ephestia*) *figulilella* (Gregson) 174
Cajanus cajan 50
Calliphoridae 98
Camellia sinensis 44
Candida chalmeris 187
Candida krusei 187
Candida oleophila 352
Carica papaya 44
Carpentaria acuminata 255
Carpophilus dimidiatus Fabricius 155
Carpophilus freemani Dobson 156
Carpophilus hemipterus (Linnaeus) 152, 155, 197
Carpophilus humeralis (Fabricius) 155
Carpophilus ligneus Murray 156
Carpophilus lugubris Murray 156
Carpophilus mutillatus Erichson 155, 156
Carpophilus obsoletus Erichson 156
Carpophilus spp. 188
Caryota mitis 238, 255
Caryota urens 238
Cassia bicapsulari 179
Casuarina tenuissima 68
Cecidomyiidae 61, 64
Cephalonomia meridionalis 147
Cephalonomia tarsalis 147
Cephalonomia waterstoni Gahan 150, 152
Cerambycidae 102
Ceratobasidiaceae 225
Ceratobasidiales 225
Ceratocystis paradoxa (Dade) 225, 227, 230
Ceratonia siliqua 179
Cerchysiella (*Zeteticontus*) *utilis* (Noyes) 155
Cermabycidae 195

- Ceroplastes floridensis* (Comstock) 332
Chalara paradoxa (De Seynes) Sacc. 225, 227
Chalara (Thielaviopsis) paradoxa 225
Chalcidoidea 64, 69
Chamaedorea elegans 238
Chamaerops humilis 21, 80, 238, 255
Chartocerus kurdjumovi Nikol'skaya 64
Cheiromenes (=*Menochilus*) *sexmaculata* (Fabricius) 34, 43, 327, 329
Cheiracanthium inclusum 341
Cheliferidae 79
Chelifer spinipalpis Redikorzev 79
Chelisoches morio (Fabricius) 98
Chelisochidae 98
Cheyletus malaccensis Oudemans 159
Chilocorus baileyi Blackburn 46
Chilocorus bipustulatus Linnaeus 34, 40, 41
Chilocorus bipustulatus var. *iranensis* 40
Chilocorus cacti L. 41
Chilocorus circumdatus (Gyllenhal) 46, 46, 327
Chilocorus nigritus (Fabricius) 43, 46
Chiroptera 309
Chondostereum purpureum 353
Chrysalidocarpus lutescens 238, 255
Chrysanthemum cinerariaefolium Vis. 358
Chrysomelidae 123
Chrysopa spp. 334
Chrysopa vulgaris Schon. 41
Chrysoperla carnea 76
Chrysoperla carnea Stephens 33, 34
Chrysoperla spp. 334
Chrysopidae 327, 41, 63
Chysalidocarpus cabadae 255
Cicadidae 66
Citrus aurantiifolia 62
Citrus aurantium 62
Citrus limon 62
Citrus maxima 62
Citrus sinensis 62
Citrus x paradisi 62
Cixiidae 70, 253
Cladosporium herbarium (Pers.Fr.) Link 234
Clubionidae 341
Coccinella septempunctata Linnaeus 34, 329
Coccinella undecimpunctata Linnaeus 34, 329, 330
Coccinellidae 57, 61, 213, 41, 63, 213, 220
Coccothrinax argentata 238
Coccothrinax argentea 255
Coccotrypes dactyliperda (Fabricius) 118
Cocos nucifera 44, 50, 82, 108, 110, 226, 238, 255
Codiaeum variegatum 54
Coffea arabica 62
Coleophoridae 72
Coleoptera 41, 57, 61, 63, 80, 102, 110, 116, 118, 120, 145, 148, 149, 151, 152, 155, 157, 158, 160, 162, 164, 122, 123, 213, 165, 213, 220
Colopalpus eriophyoides 214
Columbidae 303, 304
Columbiformes 303, 304
Comperiella bifasciata Howard 46
Comperiella lemniscata Compère & Annecke 46, 327
Coniothyrium minitans 353
Coracias benghalensis (L.) 305
Coraciidae 305
Corydalidae 334
Corypha elata 255
Corypha taliera 255
Corypha umbraculifera 110
Coxiidae 256
Crematogaster sp. 76
Criconematidae 277
Cryptolaemus montrouzieri Mulsant 57, 61
Cryptolestes ferrugineus (Stephens) 151
Cryptolestes pusillus Schönherr 149
Cucujidae 149, 151
Cunaxidae 213
Curculionidae 80
Cybocephalus nigriceps (J. Sahlbiesg) 41
Cybocephalus palmarum Peyesh. 41
Cybocephalus rutitrone Rtl. 40, 41
Cybocephalus seminulum Bau. 41
Cybocephalus tlavieeps RH. 41
Cydia pomonella Linnaeus 350, 351, 359, 360
Cyrtosus cyanipennis (Erichson) 120

D

- Dermoptera 98
Dermestidae 165
Derris elliptica 358
Deudorix isocrates 181
Deuteromycotina 115
Diaprepes abbreviates 353
Diaspididae 47

Diatreae saccharalis 351
Dichroslachys nutars 179
Dicroidiplosis manihoti Harris 327
Dicroidiplosis pseudococci Felt. 61, 64
Dictyoptera 336
Dictyosperma album 255
Dionus rubidus Motsch. 63
Diplodia phoenicum (Sacc.) H Fawc. & Klotz 235, 320
Diptera 61, 98, 109, 64, 183
Dolichodoridae 281
Dothideales 234
Drosophila melanogaster Meigen 183
Drosophilidae 183
Dufouriellus ater (Dufour) 159

E

Ectomyelois ceratoniae Zeller 177, 188, 202
Ectomyelois phoenicis (Durrant) 177
Elaeis guineensis 82, 108, 110, 238, 255
Empusa hedenborgi Stål 337
Empusa spinosa Krauss 337
Encarsia citrina Craw 46
Encyrtidae 46, 58, 61, 64
Endothia parasitica 353
Entomophthorales 220
Epeorus dowsoniella (Richard & Thompson) 175
Epeorus elutella Hübner 172
Epeorus kuehniella Zeller 202
Epeorus spp. 195, 188
Eremiaphila braueri Krauss 337
Erwinia amylovora 352
Erwinia carotovora 352
Erythraeidae 34
Eucoccidiorida 159
Eulophidae 64
Eupelmidae 69
Eupelmus sp. 69
Eupelmus sp. nr. *elongatus* Risbec 69, 327
Eurotiales 248, 249
Euryischia sp. 327
Eurytomidae 327
Euseius hibisci (Chant) 339, 340
Euseius spp. 339
Eutermes parvulus Sjostedt 140
Eutetranychus orientalis Klein 210

Exochomus flavipes Thunberg 61
Exochomus nigripennis Erichson 63

F

Ficus benghalensis (*banyan*) 44
Ficus benjamina 44
Ficus carica 44, 54, 62
Fiorinia externa Ferris 49
Fiorinia fioriniae (Targioni & Tazzetti) 47, 49
Fiorinia japonica Kuwana 49
Fiorinia juniperi (Bouche) 49
Fiorinia linderae Takagi 48
Fiorinia phoenicis Balachowsky 47
Fiorinia theae Green 49
Formicidae 76
Frankliniella schultzei (Trybom) 141
Fusarium 352
Fusarium moniliforme 353
Fusarium oxysporum 353
Fusarium oxysporum F. sp. *albedinis* Malencon 242
Fusarium oxysporum Schlechtendahl 241
Fusarium solani (Mart) sacc. 257
Fusarium spp. 352

G

Galendromus annectens (De Leon) 339
Galendromus helveolus Chant 339, 340
Galleria mellonella L. 76
Galleria mellonella Linnaeus , 202
Gelechiidae 182
Gliocladium catenulatum 353
Gliocladium virens 353
Goniozus sp. 76
Gossypium hirsutum 62
Graphiolaceae 232
Graphiolales 232
Graphiola phoenicis (Moug.) Poit. 232, 233
Gryllotalpa spp. 353
Gussia attenuate 255
Gyranusoides indica 57

H

Habrobracon hebetor Say 170
Habrocytus sp. 76

Haemogamasus pontiger 159
Hansenula fermentans 187
Haptoncus (Epuraea) luteolus (Erichson) 157
Helicosporidium parasiticum 147
Helicotylenchus dihystrera (Cobb) Sher. 280
Helicotylenchus multicinctus (Cobb) Golden 280
Helicoverpa virescens (Fabricius) 351
Helicoverpa zea (Boddie) 351
Heliothrips haemorrhoidalis Bouché 142
Hellula undalis (Fabricius) 202
Helminthosporium spp. 251, 353
Hemicronemooides mangiferae Siddiqi 277
Hemiptera 21, 98, 115, 220, 253
Hemisareoptes malus Shimer 41
Heterorhabditidae 100
Heterorhabditis bacteriophora 353
Heterorhabditis indica 101
Heterorhabditis megidis 353
Heterorhabditis sp. 100
Heterospathe elata 255
Hibiscus cannabinus 54
Hibiscus rosasinensis 54
Hibiscus sabdariffa 54
Hippodamia convergens Guérin-Méneville 333
Hippodamia quinquesignata 333
Holepyris sylvanidis (Brèthes) 147, 159
Hololena nedra 341
Homoptera 47, 50, 52, 54, 59, 62, 66, 70, 181
Hoplochelis marginalis 352
Hoplolaimidae 280
Howea belmoreana 255
Howea forsteriana 118
Hymenobosmina sp. 76
Hymenoptera 41, 46, 58, 61, 64, 76, 114, 147, 170, 178, 125, 129, 170, 202
Hyphorbe verschaffeltii 255
Hyperaspis pumila Mulsant 61, 64
Hyphaene 35
Hyphaene thebaica 42
Hyphomycetes 115
Hypoaspis sp. 104
Hypocolius ampelinus (Bonaparte) 307
Hypocreales 241, 242
Hypsicorypha gracillis Burmeister 337

I

Ichneumonidae 76, 170, 202
Isoptera 138

J

Jasminum sp. 35
Jauravia sp. 213
Jebusea hammerschmidti Reich 102, 320
Julodis euphratica Castelnau and Gory 122, 123
Julodis fimbriata Klug 122, 123
Julodis spp. 122

L

Laelapidae 104
Lasioderma serricorne Fabricius 162
Lasiodiplodia theobromae (Pat.) Grifon & Maubl 230
Latania 35
Latania sp. 255
Latrodectus bishopi Kaston 344
Latrodectus geometricus C. L. Koch 344
Latrodectus mactans (Fabricius) 344
Latrodectus variolus Walckenaer 344
Lawsonia inermis 242
Lecanicillium lecanii 350
Lepidoptera 72, 76, 77, 167, 171, 172, 174, 175, 177, 179, 182
Leptinotarsa decemlineata Say 351
Leptomastix dactylopis Howard 327, 61
Linyphiidae 343
Livistona chinensis 110, 255
Livistonia rotundifolia 255
Lobesia botrana Denis 361
Longidoridae 278, 279
Longidorus spp. 278
Lycaenidae 179
Lycosa aspersa 342
Lycosa gulosa 342
Lycosidae 342
Lymantria dispar L. 351

M

Maconellicoccus hirsutus (Green) 54, 65
Macrocentrus sp. 79

Index

- Macrocoma* sp. 123
Macroposthonia spp. 277
Macrotermes sobhyalinus Rambur 140
Makiella phoenicis K. 263
Malpighia glabra 54
Malus domestica 62
Mamestra brassicae L. 351
Mangifera indica 44, 50, 54, 62
Manilkara zapota 44, 54
Mantidae 336
Margarodidae 52
Mattesia oryzaephili 147
Mauginiella scaettae Cavara 223
Medicago sativa 242
Meloidogyne arenaria (Neal) Chitwood 274
Meloidogyne hapla (Chitwood) 274
Meloidogyne incognita (Kofoid & White) Chitwood 274
Meloidogyne javanica (Treub) Chitwood 274
Meloidogynidae 274
Melolontha melolontha 352
Melyridae 120
Metarhizium anisopliae (Metchnikoff) 115, 350, 352
Metarhizium anisopliae var. *acridium* 135, 136
Metarhizium flavoviride 352
Metaseiulus occidentalis (Nesbitt) 325
Metroxylon sagu 82
Mexecheles hawaiiensis (Baker) 217
Microcerotermes buettikeri Chhotani & Bose 140
Microcerotermes diversus Silvestri 138, 140
Microphthalma disjuncta (Wiedemann) 109
Microsporidia 159
Microthrips dmitriewi Werner. 337
Misumena vatia 343
Monilliaceae 223
Monilliales 223
Mucoraceae 250
Mucorales 250
Muridae 290, 291, 293
Musa sp. 44
Mus musculus (Linnaeus) 286, 291
Mycetophagidae 164
Mycosphaerellaceae 234
Mycosphaerella tassiana (De Not.) Johans 234
Mycterotermes meringocephalus Sands 140
Myelois ceratoniae Zeller 177
Myelois phoenicis Durrant 177
Myndus crudus Van Duzee 253, 256
Myrmeleotidae 334
Myrothecium verrucaria 353
- N**
- Nannorrhops ritchiana* 255
Nectriaceae 241, 242
Nemeritis canescens Gravenhorst 170
Nemeritis (Venturia) canescens Gravenhorst 202
Neochrysocharis formosa (Westwood) 64
Neodyspis decaryi 255
Neoseiulus californicus Mc Gregor 325, 339, 340
Neoseiulus fallacis Garman 219
Neowashingtonia 35
Neozygitaceae 220
Neozygites adjarica 220
Nephus quadrimaenatus Hbst. 41
Neuroptera 41, 63
Nicotiana rustica L. 356
Nipaecoccus vastator (Maskell) 62
Nipaecoccus viridis (Newstead) 62, 65
Nitidulidae 41, 152, 155, 157
Nosema locustae 353
Nosema oryzaephili 147
Nosematidae 159
Nosema whitei 147, 159
- O**
- Olea europaea* L. 50
Oliarus atkinsoni 70
Oligonychus afrasiaticus (McGregor) 205
Oligonychus pratensis Banks 217, 220
Oligonychus sacchari 209
Ommatissus binotatus 21
Ommatissus binotatus var. *lybicus* 21
Ommatissus lybicus Bergevin 22, 21
Omphalia pigmentata Bliss 245
Omphalia spp. 257
Omphalia tralucida Bliss 245
Oncophylla samadina 325
Orius albidipennis Reuter 159
Orius laevigatus (Fieber) 159
Orius spp. 220
Orthoptera 131, 136
Oryctes agamemnon Burmeister 105
Oryctes boas Fabricius 108
Oryctes elegans Prell. 107, 108

- Oryctes monoceros* (Olivier) 109
Oryctes rhinoceros (Linnaeus) 98, 110
Oryctes sahariensis De Mire 109
Oryctes spp. 105, 109
Oryzaephilus mercator (Fauvel) 148
Oryzaephilus surinamensis (Linnaeus) 145, 205
Oryza sativa 151
Ostrinia nubilalis (Hubner) 359
Ostrinia spp. 352
Otiorhynchus sulcatus 353
Otiorhynchus sulcatus (F.) 326
- P**
- Pachnaeus litus* 353
Paecilomyces fumosoroseus 352
Panonychus ulmi (Koch) 219
Paralongidorus spp. 278
Parasierola sp. 76
Parlatoria blanchardi (Targioni- Tozetti) 35, 42, 349
Passer domesticus (L.) 302
Passeridae 302
Passeriformes 302, 305, 306, 307
Pasteurella pestis 289
Pediculoides ventricosus (Newport) 202
Penicillium spp. 187, 249
Peregrinator biannulipes Montrouzier & Signoret 159
Perindus binudatus Emeljanov 70, 256
Peronosporales 239
Persea americana 54
Pestalotiopsis palmarum (Cook) Steyaert 237
Phanerotoma flavitestacea Fischer , 178, 202
Phanerotoma ocularis Kohl. 76, 79
Pharoscymnus anchorago (Fairmaire) 41, 43
Pharoscymnus flexibilis Mulsant 41, 43
Pharoscymnus horni (Weise) 40, 41, 43
Pharoscymnus numidicus Pic. 40, 41, 41
Pharoscymnus smirnovi Dobzha. 41
Pharoscymnus tristiculus Sicard 327
Phidippus otiosus 342
Phoenicococcus marlatti Cockerell 42
Phoenix 35
Phoenix canariensis 42, 238, 255
Phoenix dactylifera 44, 54, 62, 80, 108, 110, 179, 238, 255
Phoenix reclinata 238, 255
Phoenix roebelenii 238, 255
Phoenix sylvestris 238, 255
- Phoenix theophrasti* 238
Phomopsis spp. 352
Phonopate frontalis Fåhraeus 120
Photorhabdus 100
Photorhabdus spp. 326
Phytonemus pallidus (Banks) 219
Phytophthora palmivora 320
Phytophthora sp. nr. *palmivora* 239
Phytoseiidae 213, 219
Phytoseiulus macropilis (Banks) 216
Phytoseiulus persimilis Athias-Henriot 325, 340
Phytoseiulus spp. 339
Planococcus citri (Risso) 59, 65
Platynemeris laevicollis Distant 98, 115
Platyleura arabica Myers 66
Pleosporaceae 247, 251
Pleosporales 247, 251
Plodia interpunctella (Hübner) 147, 170, 175, 188, 202
Plutella xylostella 351
Polistes annularis L. 130
Polistes exclamans Vier. 130
Polistes foederatus Kol. 130
Polistes fuscatus Cr. 130
Polistes gallicus (Linnaeus) 129
Polistes hebraeus F. 130
Polistes nymphus Chr. 130
Prays oleae Bern 345
Pritchardia 35
Pritchardia affinis 255
Prochiloneurus aegyptiacus (Mercet) 64
Prosopis piralis 179
Prosopis spicigera 132
Pseudaspidioproctus hyphaeniacus Hall. 52
Pseuderimerus sp. , 69
Pseudococcidae 54, 59, 62
Pseudoligosita babylonica Viggiani 34, 327
Pseudomonas chloraphis 352
Pseudomonas fluorescens 352
Pseudomonas lachrymans Carsner 356
Pseudomonas syringae 352
Pseudomonas tabaci Stevens 356
Pseudophilus testaceus (Gah.) 102
Psidium guajava 44, 54
Pteromalidae 76
Pteropodidae 309
Ptychosperma macarthurii 255
Pulicidae 288

Pullus ebner Wse. 41
Punica granatum 62, 179
Punica vulgaris 179
Pycnonotidae 306
Pycnonotus xanthopygos (Gould) 306
Pyemotes herfsi (Oudemans) 170, 202
Pyemotidae 170, 202
Pyratidae 76, 77, 167, 171, 172, 174, 175, 177
Pyrus communis 62
Pythiaceae 239
Pythium oligandrum 353
Pythium spp. 352

R

Raoiella indica Hirst 211, 213
Raphia ruffia 110
Rattus norvegicus (Berkenhout) 286, 293
Rattus rattus (Linnaeus) 286, 290
Ravenea hildebrandtii 255
Reduviidae 98, 115
Rhabdionvirus oryctes Hüger 113
Rhipidophyllum hystrix 255
Rhapis excelsa 238
Rhizoctonia spp. 352, 353
Rhizopus nigricans Ehrenberg 250
Rhynchophorus bilineatus (Montrouzier) 82
Rhynchophorus cruentatus (Fabricius) 82
Rhynchophorus ferrugineus (Olivier) 80, 87, 362
Rhynchophorus palmarum Linnaeus 82
Rhynchophorus papuanus Kirsch 82
Rhynchophorus phoenicis Fabricius 81, 82
Rhynchophorus schach Olivier 82
Rhynchophorus spp. 320
Rhynchophorus vulneratus (Panzer) 82
Rodentia 290, 291, 293
Rodolia (Vedalia) cardinalis Mulsant 61
Rousettus aegyptiacus (Geophroy) 309
Roystonea elata 238, 255
Roystonea hispaniola 255
Roystonea regia 255
Runcinia sp. 34, 34
Ryania speciosa Vahl. 359

S

Sabal causiarum 255
Sabal minor 238

Sabal palmetto 82, 238, 255
Saccharopolyspora spinosa 355
Saccharum officinarum 82
Saccnaromyces cervesiae 187
Salticidae 342
Sarcophaga fuscicauda Böttcher 98
Sascoptidae 41
Scapteriscus vicinus 353
Scarabaeidae 110, 116
Schistocerca gregaria Forskål 131
Scolia oryctophaga Coquerel 114
Scolia ruficornis Fabricius 114
Scoliidae 114
Scolothrips sexmaculatus (Pergande) 219
Scolytidae 118
Scymnus levaillanti Mulsant 327
Scymnus sp. 329
Sialidae 334
Silvanidae 145, 148
Silvanus spp. 195
Siphonaptera 288
Sitotroga cerealella (Olivier) 182
Solanum melongena 50
Sorghum bicolor 151
Spectrobates *ceratoniae* Zeller 177
Sphodromantis trimacula Saussure 337
Spirillum minus Sodoku 288
Spodoptera exigua 351
Spodoptera exigua (Hubner) 351
Spodoptera spp. 351
Steinernema abbasi 101
Steinernema carpocapsae (weiser) 100, 353
Steinernema feltiae 350, 353
Steinernema glaseri 353
Steinernema riobravae 101, 100
Steinernema scapterisci 353
Steinernema spp. 326
Steinernematidae 100
Stemphylium sp. 251
Stethorus darwini (Brèthes) 332
Stethorus exspectatus Chazeau 332
Stethorus gilvifrons (Mulsant) 209, 332
Stethorus histrio Chazeau 332
Stethorus japonicus Kamiya 332
Stethorus keralicus Kapur 213, 213
Stethorus loi Sasaji 332
Stethorus loxtoni Britton & Lee 332

- Stethorus madecassus* Chazeau 332
Stethorus parcempunctatus 213
Stethorus pauperculus (Weise) 332
Stethorus punctillum (Weise) 209, 332
Stethorus punctum (Leconte) 220, 332
Stethorus spp. 332
Stethorus tetranychi Kapur 213
Stethorus tridens Gordon 332
Sticholotis punctata Crotch 217
Stigmeidae 338
Stomella radicicola 320
Strategus julianus Burmeister 116
Streptobacillus moniliformis 288
Streptomyces avermitilis 354
Streptomyces griseoviridis 352
Streptomyces griseus 356
Streptomyces hygroscopicus subsp. *aureolacrimosus* 355
Streptopelia senegalensis (L.) 303
Streptopelia tutur (L.) 304
Streptoverticillium rimofaciens 355
Syagrus amara 255
Syagrus romanzoffiana 238, 255
Syagrus schizophylla 255
- T**
- Tachinidae 109
Tamarindus indica 44
Tanacetum cinerariaefolium 358
Telsimia ephippiger Chapin 213
Tenebrionidae 158, 160
Tenuipalpidae 211, 214, 215
Tenuipalpus eriophyoides Baker 214
Termatophyllum insigne Reuter 159
Termitidae 138
Tetranychidae 205, 210, 217
Tetranychus spp. 219
Tetranychus urticae 219, 220
Tetranychus urticae Koch. 361
Theobroma cacao 50
Theridiidae 343
Theridion dilutum 343
Theridion fordum 343
Thielaviopsis paradoxa (De Seynes) Höhn 225, 227, 230
Thomisidae 327, 343
Thrinax morrisii 255
- Thrinax parviflora* 255
Thripidae 141, 142, 142, 219
Thysanoptera 141, 142, 142, 219
Torulopsis dactylifera 187
Torulopsis pulcherrina 187
Torymidae 69
Trachycarpus fortunei 255
Trialeurodes vaporariorum 352
Tribolium castaneum Herbst 160
Tribolium confusum Jacquelin du Val 158
Tribolium spp. 195
Trichocomaceae 248, 249
Trichoderma harzianum 353
Trichogramma brassicae Bezdenko 181, 327
Trichogramma cacoeciae Marchal 346
Trichogramma chilonis Ishii (= *T. confusum* Viggiani) 346
Trichogramma chilotraeae 181
Trichogramma dendrolimi Matsumura 346
Trichogramma evanescens 346
Trichogramma evanescens (Westwood) 346
Trichogramma japonicum Ashmead 346
Trichogramma maidis Pintureau & Voegelè 327, 346
Trichogramma minutum Riley 346
Trichogramma oleae Voegelè & Pointel 346
Trichogramma ostriniae Pang & Chan 346
Trichogramma platneri Nagarkatti 346
Trichogramma pretiosum Riley 346
Trichogramma semblidis (Aurivillius) 346
Trichogramma spp. 76, 181, 182
Trichogrammatidae 181
Tricholomataceae 245
Trogoderma granarium Everts 165, 320
Tropiduchidae 21
Tylenchida 274, 276, 277, 280, 281
Tylenchorhynchus spp. 281
Typhaea stercorea (Linnaeus) 164
Typhlodromus caudatus Chant 213
Tyrophagus lintneri Osborne 185
Tyrophagus putrescentiae (Schrank) 185
- U**
- Ulocadium atrum* Presuss 231
Unaspis citri (Comstock) 332

V

- Veitchia merrillii* 238
Veitchia spp. 255
Verticillium lecanii 352
Vespa orientalis L. 125
Vespidae 125, 129
Vicia faba 179
Vinca sp. 35
Virachola (Deudorix) livia (Klug) 179, 345
Virachola livia Klug 202
Vitis vinifera 62

W

- Washingtonia* 35
Washingtonia filifera 42, 80, 255
Washingtonia robusta 238, 255
Washingtonia spp. 108, 238

X

- Xanthomonas oryzae* Dows. 356
Xanthomonas citri Dows. 356
Xenopsylla cheopis (Rothschild) 288, 289
Xenorhabdus 100
Xenorhabdus spp. 100
Xiphinema americanum Cobb 279
Xylariales 237
Xylocoris cursitans 147
Xylocoris flavipes Reuter 147, 159
Xylocoris sordidus (Reuter) 159

Y

- Yersinia pestis* 289

Z

- Zea mays* 151
Ziziphus mauritiana 62
Zygosaccharomyces barkeri 187
Zygosaccharomyces cavarae 187
Zygosaccharomyces globiformis 187